



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A systematic review of associations between maternal exposures during pregnancy other than smoking and antenatal fetal measurements

Citation for published version:

Huang, I, Mak, D, Cheung, P, Abraham, M, Clemens, T & Turner, S 2019, 'A systematic review of associations between maternal exposures during pregnancy other than smoking and antenatal fetal measurements', *Environmental Research*, vol. 173, pp. 528-538.
<https://doi.org/10.1016/j.envres.2019.04.005>

Digital Object Identifier (DOI):

[10.1016/j.envres.2019.04.005](https://doi.org/10.1016/j.envres.2019.04.005)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Environmental Research

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Manuscript Number: ER-18-2630R1

Title: A systematic review of associations between maternal exposures during pregnancy other than smoking and antenatal fetal measurements

Article Type: Research paper

Section/Category: Exposure

Keywords: Air pollution; Benzene; Diet; Ethanol; Fetus; Mother; Phthalic acids.

Corresponding Author: Dr. Steve William Turner, MD

Corresponding Author's Institution: University of Aberdeen

First Author: Ivory Huang

Order of Authors: Ivory Huang; Diane Mak; Prudence Cheung; Miriam Abraham; Tom Clemens; Stephen William Turner, MD

Abstract: Background. Reduced birth weight is associated with many maternal environmental exposures during pregnancy, but the gestational age at onset of this association is unknown. We have previously reported associations between maternal smoking and fetal size.

Objective. To report on our systematic review of the literature describing associations between antenatal size and growth and maternal exposures during pregnancy.

Data sources. Electronic databases (OVID and EMBASE) and web sites for cohort studies were searched.

Study eligibility. Studies were eligible if they examined associations between maternal environmental exposures (including ambient air exposure, diet and alcohol) and antenatal fetal ultrasound measurements.

Study appraisal. The Navigation Guide was used to assess the strength of evidence.

Results. There were 451 abstracts identified and 36 papers were included of which maternal diet was the exposure of interest in 15, maternal ambient air exposure in 10, maternal alcohol in 3 and other exposures in 8. The first paper was published in 2006. Associations were present between exposures in 18% of comparisons with second trimester measurements and in 46% of comparisons with third trimester measurements. In the third trimester, when an association was present, reduced head size was most commonly (58%) associated with current or previous maternal exposure, with reduced length being least commonly (32%) associated and reduced weight being intermediate (52%). In the third trimester, increased maternal nitrogen dioxide exposure was associated with reduced head size was associated with in all seven studies identified and reduced fetal weight in five out of six studies.

Conclusion. There is sufficient evidence of toxicity in the context of maternal exposure to nitrogen dioxide and reduced third trimester fetal head size. There is insufficient evidence of toxicity with regard to maternal exposures to dietary factors, alcohol and environmental chemicals and reduced fetal size.

Response to Reviewers: Thank you very much for the opportunity to submit a revision of our manuscript. Thank you also to the reviewer for their time and helpful comments. We have amended the manuscript in light of the reviewer's comments (see point by point response below). We have also included a "Highlights" file in the submission. Finally, we have identified a paper missed in our earlier manuscript (Lopez-Espinosa 2016) and have included this in the revised manuscript.

With kind regards

Steve Turner (on behalf of all authors)

Reviewer #2: The manuscript reviewed literature of dietary factors, ambient air pollution, alcohol intake, and other environmental factors and fetal growth assessed during pregnancy using ultrasound. The number of studies on each of the exposures was relatively small for most exposures, which significantly limited the ability to provide more definitive answers to these associations. Nevertheless, the findings on NO₂ and PM_{2.5} were more consistent compared with other exposures. Overall, the manuscript was well prepared, and the coverage of study was excellent. The manuscript provided quality of the different aspects of the study, and the sample size of many studies are large.

While the findings are considered reliable and faithful to the original studies, the review has several limitations.

1. The exposures with less than 3 studies are not the best case for systematic review, as the limited studies will reduce the credibility of the review.

OUR RESPONSE. We accept that this is a limitation of the literature, but not of our review. We have replaced the sentence in the limitations section of the discussion in the original submission and emphasised this important point.

2. The exposure and ultrasound measurement were different for timing during pregnancy, the comparability between different studies is a concern. Fetal growth on a population level may be easy to describe by gestational age, the individual difference cannot be dismissed.

OUR RESPONSE. The reviewer is right that fetal measurements and exposures were determined at different gestational ages between and also within studies. We have added to the limitations section of the discussion to acknowledge this point.

3. It was not clear how maternal conditions were handled in many studies included. The representativeness of the study participants was a concern by the authors themselves, as the risk of bias on recruitment was flagged.

OUR RESPONSE. As the reviewer states, we highlight the bias in recruitment strategy to the majority of the papers included in the review. The studies are not able to address/adjust for risk of bias in recruitment in their analyses. We think that by "maternal conditions" the reviewer is referring to covariates such as maternal smoking status, age, socioeconomic status, etc. In "Determining risk of bias" (see the supplement) we explain how our methodology considered (i) whether maternal smoking status was considered as an essential covariate (ii) whether the analysis included adjustment for covariates and (iii) whether covariates were described and were comprehensive.

4. Covariates adjustment is important for an individual study, but on the systematic review's standpoint, the difference in covariates adjustment may not be trivial.

OUR RESPONSE. The reviewer makes a valid observation does not ask us to amend the manuscript. We agree that for each separate study, the effect of each covariate on the outcome (i.e. fetal growth) will likely be different. If we had undertaken meta analysis using unadjusted data this could have been an important consideration, but this issue is not as relevant to our systematic review.

5. Consistency with birth weight (or small for gestational age) was not provided while literature in that regard is more abundant

OUR RESPONSE. Thank you for this helpful suggestion. We did mention the literature linking maternal exposures to birth weight in the introduction of the original manuscript. We have added a new paragraph in the discussion which considers the consistency of the literature describing associations between maternal exposure and birth weight and fetal size.

Research Data Related to this Submission

There are no linked research data sets for this submission. The following reason is given:

The data are already published in the literature. Original data requests should be directed to the authors of the original papers cited in our review.



Child Health
 Royal Aberdeen Children's Hospital
 Aberdeen
 AB25 2ZG
 30th Mar 2019

Dear Editor,

Thank you very much for the opportunity to submit a revision of our manuscript. Thank you also to the reviewer for their time and helpful comments. We have amended the manuscript in light of the reviewer's comments (see point by point response below). We have also included a "Highlights" file in the submission. Finally, we have identified a paper missed in our earlier manuscript (Lopez-Espinosa 2016) and have included this in the revised manuscript.

With kind regards

Steve Turner

Reviewer #2: The manuscript reviewed literature of dietary factors, ambient air pollution, alcohol intake, and other environmental factors and fetal growth assessed during pregnancy using ultrasound. The number of studies on each of the exposures was relative small for most exposures, which significantly limited the ability to provide more definitive answers to these associations. Nevertheless, the findings on NO₂ and PM_{2.5} were more consistent compared with other exposures. Overall, the manuscript was well prepared, and the coverage of study was excellent. The manuscript provided quality of the different aspects of the study, and the sample size of many studies are large.

While the findings are considered reliable and faithful to the original studies, the review has several limitations.

1. The exposures with less than 3 studies are not the best case for systematic review, as the limited studies will reduce the credibility of the review.
 OUR RESPONSE. We accept that this is a limitation of the literature, but not of our review. We have replaced the sentence in the limitations section of the discussion in the original submission and emphasised this important point.
2. The exposure and ultrasound measurement were different for timing during pregnant, the comparability between different studies is a concern. Fetal growth on a population level may be easy to describe by gestational age, the individual difference cannot be dismissed.
 OUR RESPONSE. The reviewer is right that fetal measurements and exposures were determined at different gestational ages between and also within studies. We have added to the limitations section of the discussion to acknowledge this point.
3. It was not clear how maternal conditions were handled in many studies included. The representativeness of the study participants was a concern by the authors themselves, as the risk of bias on recruitment was flagged.
 OUR RESPONSE. As the reviewer states, we highlight the bias in recruitment strategy to the majority of the papers included in the review. The studies are not able to address/adjust for risk of bias in recruitment in their analyses. We think that by "maternal conditions" the reviewer is referring to covariates such as maternal smoking status, age, socioeconomic status, etc. In "Determining risk of bias" (see the

supplement) we explain how our methodology considered (i) whether maternal smoking status was considered as an essential covariate (ii) whether the analysis included adjustment for covariates and (iii) whether covariates were described and were comprehensive.

4. Covariates adjustment is important for an individual study, but on the systematic review's standpoint, the difference in covariates adjustment may not be trivial.

OUR RESPONSE. The reviewer makes a valid observation does not ask us to amend the manuscript. We agree that for each separate study, the effect of each covariate on the outcome (i.e. fetal growth) will likely be different. If we had undertaken meta analysis using unadjusted data this could have been an important consideration, but this issue is not as relevant to our systematic review.

5. Consistency with birth weight (or small for gestational age) was not provided while literature in that regard is more abundant

OUR RESPONSE. Thank you for this helpful suggestion. We did mention the literature linking maternal exposures to birth weight in the introduction of the original manuscript. We have added a new paragraph in the discussion which considers the consistency of the literature describing associations between maternal exposure and birth weight and fetal size.

Thank you very much for the opportunity to submit a revision of our manuscript. Thank you also to the reviewer for their time and helpful comments. We have amended the manuscript in light of the reviewer's comments (see point by point response below). We have also included a "Highlights" file in the submission. Finally, we have identified a paper missed in our earlier manuscript (Lopez-Espinosa 2016) and have included this in the revised manuscript.

With kind regards

Steve Turner (on behalf of all authors)

Reviewer #2: The manuscript reviewed literature of dietary factors, ambient air pollution, alcohol intake, and other environmental factors and fetal growth assessed during pregnancy using ultrasound. The number of studies on each of the exposure was relative small for most exposures, which significantly limited the ability to provide more definitive answers to these associations. Nevertheless, the findings on NO₂ and PM_{2.5} were more consistent compared with other exposures. Overall, the manuscript was well prepared, and the coverage of study was excellent. The manuscript provided quality of the different aspects of the study, and the sample size of many studies are large.

While the findings are considered reliable and faithful to the original studies, the review has several limitations.

1. The exposures with less than 3 studies are not the best case for systematic review, as the limited studies will reduce the credibility of the review.
OUR RESPONSE. We accept that this is a limitation of the literature, but not of our review. We have replaced the sentence in the limitations section of the discussion in the original submission and emphasised this important point.
2. The exposure and ultrasound measurement were different for timing during pregnant, the comparability between different studies is a concern. Fetal growth on a population level may be easy to describe by gestational age, the individual difference cannot be dismissed.
OUR RESPONSE. The reviewer is right that fetal measurements and exposures were determined at different gestational ages between and also within studies. We have added to the limitations section of the discussion to acknowledge this point.
3. It was not clear how maternal conditions were handled in many studies included. The representativeness of the study participants was a concern by the authors themselves, as the risk of bias on recruitment was flagged.
OUR RESPONSE. As the reviewer states, we highlight the bias in recruitment strategy to the majority of the papers included in the review. The studies are not able to address/adjust for risk of bias in recruitment in their analyses. We think that by "maternal conditions" the reviewer is referring to covariates such as maternal smoking status, age, socioeconomic status, etc. In "Determining risk of bias" (see the supplement) we explain how our methodology considered (i) whether maternal smoking status was considered as an essential covariate (ii) whether the analysis included adjustment for covariates and (iii) whether covariates were described and were comprehensive.
4. Covariates adjustment is important for an individual study, but on the systematic review's standpoint, the difference in covariates adjustment may not be trivial.
OUR RESPONSE. The reviewer makes a valid observation does not ask us to amend the manuscript. We agree that for each separate study, the effect of each covariate on the outcome (i.e. fetal growth) will likely be different. If we had undertaken meta analysis

using unadjusted data this could have been an important consideration, but this issue is not as relevant to our systematic review.

5. Consistency with birth weight (or small for gestational age) was not provided while literature in that regard is more abundant
OUR RESPONSE. Thank you for this helpful suggestion. We did mention the literature linking maternal exposures to birth weight in the introduction of the original manuscript. We have added a new paragraph in the discussion which considers the consistency of the literature describing associations between maternal exposure and birth weight and fetal size.

*Highlights

- Many modifiable maternal exposures are associated with reduced birth weight
- Here we ask the question are maternal exposures linked with reduced fetal size?
- Maternal exposure to air pollutants is inversely associated with fetal head size
- It is not clear whether other maternal exposures are important to fetal size
- There is a need for more research into determinants of fetal size

ABSTRACT

Background. Reduced birth weight is associated with many maternal environmental exposures during pregnancy, but the gestational age at onset of this association is unknown.

We have previously reported associations between maternal smoking and fetal size.

Objective. To report on our systematic review of the literature describing associations between antenatal size and growth and maternal exposures during pregnancy.

Data sources. Electronic databases (OVID and EMBASE) and web sites for cohort studies were searched.

Study eligibility. Studies were eligible if they examined associations between maternal environmental exposures (including ambient air exposure, diet and alcohol) and antenatal fetal ultrasound measurements.

Study appraisal. The Navigation Guide was used to assess the strength of evidence.

Results. There were 451 abstracts identified and 365 papers were included of which maternal diet was the exposure of interest in 15, maternal ambient air exposure in 10, maternal alcohol in 3 and other exposures in 87. The first paper was published in 2006.

Associations were present between exposures in 189% of comparisons with second trimester measurements and in 464% of comparisons with third trimester measurements. In the third trimester, when an association was present, reduced head size was most commonly (5869%) associated with current or previous maternal exposure, with reduced length being least commonly (3227%) associated and reduced weight being intermediate (5246%). In the third trimester, increased maternal nitrogen dioxide exposure was associated with reduced head size was associated with in all seven studies identified and reduced fetal weight in five out of six studies.

Conclusion. There is sufficient evidence of toxicity in the context of maternal exposure to nitrogen dioxide and reduced third trimester fetal head size. There is insufficient evidence of toxicity with regard to maternal exposures to dietary factors, alcohol and environmental chemicals and reduced fetal size.

Key words. Air pollution; Benzene; Diet; Ethanol; Fetus; Mother; Phthalic acids

ABBREVIATIONS

HC=head circumference

AC=abdominal circumference

BPD=biparietal diameter

BTEX= aromatic hydrocarbons (benzene, toluene, ethylbenzene, m/p-xylene, and o-xylene)

CO=carbon monoxide

4,4'DDE=4,4'dichlorodiphenylchloroethylene

EFW=estimated fetal weight.

HCB= hexachlorobenzene

MAD=mean abdominal circumference

NCD=non-communicable diseases

NO₂= nitrogen dioxide

O₃=ozone

PAH=polycyclic aromatic hydrocarbons

PCB=polychlorinated biphenyls

PM₁₀=particulates with diameter less than 10 microns

PUFA=poly unsaturated fatty acids

SO₂=sulphur dioxide

INTRODUCTION

Being small for gestational age (SGA) at birth is associated with increased risk for conditions that include coronary artery disease(Barker 1995a), type II diabetes(Hales CN and Barker DJ 1992) and asthma(Shaheen SO et al. 1999). The fetal origins hypothesis(Barker 1995a) and more recently, concepts including developmental plasticity(Bateson et al. 2004) and predictive adaptive responses(Gluckman et al. 2005), suggest that some antenatal exposures (or cues) predispose an unborn individual to non-communicable diseases (NCD) in later life by a mechanism which involves reduced fetal size and growth. A recent systematic review linking reduced fetal size, as evidenced by fetal ultrasound scan, to NCD in children supports the principle of the developmental origins of disease(Alkandari et al. 2015).

The mechanism(s) where antenatal cues reduce fetal size and pre-dispose to NCD are not fully understood, but are thought to include epigenetic modification in fetal cells following maternal exposure during pregnancy(Gluckman et al. 2011). Maternal exposures linked to SGA at birth include maternal smoking(Pereira et al. 2017), exposures to poor quality ambient air(Maisonet et al. 2004; Shah PS. Balkhair T. Knowledge Synthesis Group on Determinants of Preterm/LBW births 2011), dietary factors(Gresham E et al. 2014) and exposure to chemicals such as bisphenol A(Perera F and Herbstman J 2011). Knowledge of which exposures are associated with reduced antenatal size and the gestation at which these exposures may be acting would be important to our understanding of “fetal origins” and for public health educational messages and interventions, but the literature describing associations between these exposures and fetal size has not been systematically reviewed. Our group has recently undertaken a systematic review of the literature linking antenatal size and growth to maternal smoking and found that maternal smoking is consistently associated with fetal growth failure after the second trimester(Abraham et al. 2017). Here, we undertake a systematic review designed to answer the population exposure, comparator and outcome (PECO) question: Are fetuses who are exposed to maternal environmental exposures other than smoking small for gestational age compared to unexposed fetuses?

The Navigation Guide methodology is ideally suited for determining the strength of evidence between exposures and fetal size (Johnson et al. 2014) and was used in our study.

METHODS

Search methodology

A database search was carried out September 2017 using OVID MEDLINE and Embase databases and updated in May 2018. Search terms were developed initially using those used in previous systematic reviews (Abraham et al. 2017; Alkandari et al. 2015) and modified after identifying relevant publications not identified using these previous search terms (Aguilera I et al. 2010; Slama R et al. 2009). The online supplement shows the search terms used. Additional papers were identified from bibliographies and from the web sites of the following cohorts known to have collected ultrasound measurements of fetal size: the Raine cohort (<http://www.rainestudy.org.au/>), the EDEN cohort (<https://eden.vjf.inserm.fr>), Southampton Women Study (SWS, <http://www.leu.soton.ac.uk/sws/>), Generation R (<http://www.erasmusmc.nl/epi/research/Generation-R/>), and INMA Mother and Child Cohort Study (Guxens M et al. 2012). The supplement describes these cohorts in greater detail. Two researchers (MA and ST or IH and ST) independently reviewed abstracts identified by the database search and full papers considered potentially eligible were obtained. Eligible papers had to report fetal ultrasound anthropometric measurements (i.e. crown rump length, biparietal diameter, head circumference, femur length, abdominal circumference or estimated fetal weight) and relate these to a current or previous maternal environmental exposures. From background reading, the following exposures were sought: ambient air pollution, maternal alcohol ingestion, maternal diet, maternal drug use, occupational exposures, or pesticide exposure. Papers where fetal size was related to obstetric complications, maternal smoking (as the primary exposure) and maternal drug treatment were excluded. The outcome measures were fetal size and fetal growth.

Fetal measurements. First trimester (i.e. ≤ 13 weeks gestation) measurements were: crown rump length (CRL), biparietal diameter (BPD), head circumference (HC) abdominal

circumference (AC) and mean abdominal diameter (MAD). Second trimester (i.e. >13 to <28 weeks gestation) and third trimester (i.e. ≥28 weeks) measurements were femur length (FL), HC, BPD, MAD, AC and estimated fetal weight (EFW) (the latter was derived from BPD, AC and FL measurements (Hadlock et al. 1984)).

Quality assessment of data

The Navigation Guide methodology was used to assess the risk of bias and the quality and strength of evidence (Johnson et al. 2014). We developed a protocol to assess risk of bias, quality of evidence and strength of evidence and which was based on the study of Johnson *et al* (Johnson et al. 2014) and which is described in the supplement. Risk of bias was rated for each paper and then across each of four exposure groups (i.e. maternal dietary exposures, air pollution exposures, alcohol exposure and other exposures) to inform the rating of quality of the evidence. The quality of the evidence was then used as part of the assessment of the strength of evidence. Two researchers (DM and PC) independently assessed each paper and then consensus was reached during discussion with three researchers (DM, PC and ST).

Meta-Analysis of data

We sought to identify results which might be suitable for meta-analysis. To do this we identified exposures where the same units of exposure were used in studies using comparable methodologies and reporting the same fetal measurement outcome.

RESULTS

Papers identified

The search identified 451 abstracts and 365 papers were ultimately included in this review, (figure 1). Fifteen papers considered maternal dietary factors (Bergen et al. 2016; Bouwland-Both MI et al. 2013; Carlsen K et al. 2013; Drouillet P et al. 2009; Drouillet-Pinard P et al. 2010; Heppe DH et al. 2011a; Heppe DH et al. 2011b; Ioannou C et al. 2012; Karateke et al.

2015; Mahon P et al. 2010; Steenweg-de Graaff et al. 2017; Timmermans S et al. 2009; Timmermans S et al. 2012; Turner SW et al. 2010; Young BE et al. 2012), ten ambient air exposures(Aguilera I et al. 2010; Carvalho M.A. et al. 2016; Clemens et al. 2017; Hansen CA et al. 2008; Iniguez C et al. 2012; Iniguez et al. 2016; Malmqvist et al. 2017; Ritz B et al. 2014; Slama R et al. 2009; van den Hooven EH et al. 2012), three maternal alcohol consumption(Bakker R et al. 2010; Handmaker NS et al. 2006; Kfir M et al. 2009) and eightseven considered other exposures(Botton et al. 2016; Casas et al. 2016; Harari et al. 2015; Lopez-Espinosa et al. 2015; Philippat et al. 2014; Snijder CA et al. 2012b; Snijder CA et al. 2013; Lopez-Espinosa et al. 2016). Studies which related environmental exposures to fetal anomalies (e.g. kidney cysts) and one relating physical demands of work(Snijder CA et al. 2012a) to fetal size were excluded. All studies were observational and the first was published in 2006. Three publications were from the USA, one each from Australia, Argentina, Brazil, Pakistan and Ukraine and the remaining 287 from European populations. Twenty one publications were from one of three birth cohorts (including ten from the Generation R cohort, sixfive from INMA and five from EDEN).

Risk of bias

There was generally a low risk of overall bias across the studies, with the exception of recruitment bias and to a lesser extent, exposure assessment, confounding and incomplete outcome data. Recruitment bias was low in three studies, probably low in 298 studies and high in another four, figure two. Risk of bias in exposure assessment was low in 176 studies, probably low in 8 and probably high in 11, figure two. Bias due to confounding was low risk in 30, probably low risk in onetwo, probably high risk in a further fourtwo and high risk in one study. Incomplete outcome data was at low risk for bias in 18, probably low risk in ten, probably high risk in seven and high risk in one study. Risk for bias from selective reporting was low in 332 studies, probably low in two more and probably high in one. Supplemental table one explains reasons why a study was not rated at low risk of bias for the eight

domains considered. The quality and strength of evidence are described later for each separate category of exposure.

Overview of findings

Table 1 describes the exposure, the trimester where an association was sought and the direction of any association between exposures and fetal measurement. Regarding consistency of findings for second trimester measurements, an association was present between a maternal exposure in 16 of 8~~84~~ (1~~89~~%) comparisons made, and these associations were distributed evenly across all fetal measurements and all the exposures measured (table 1 and supplemental table 2). Third trimester measurements were associated with maternal exposures in 44~~34~~ of the 9~~63~~ comparisons made (4~~64~~%) and were present for 15 of 2~~65~~ (5~~860~~%) analyses where BPD or HC were considered, 9~~7~~ of 2~~86~~ (3~~227~~%) where FL was considered, 8 of 1~~98~~ (4~~24~~%) where AC was considered and 1~~24~~ of 2~~32~~ (5~~20~~%) where EFW was considered (table 1 and supplemental table 2). Meta-analysis was not possible for maternal exposures to dietary factors, alcohol and other chemicals since either only one exposure was reported in the literature or if more than one study reported the same outcome, it was not valid or possible to pool the data. Meta-analysis was therefore only considered for ambient air exposures since comparable data seemed available (but not for other exposure categories), however the studies identified used a variety of methods to measure and report exposure (including units) meaning that meta analysis was not possible (table 2).

Maternal dietary exposures

Maternal plasma nutrient concentrations and fetal measurements

Vitamin D. FL was linked to maternal plasma 25(OH)D hydroxyl vitamin D in one study of 171 mothers aged <18 years (Young BE et al. 2012) and reduced exposure (i.e. ≤ 50 nmol/L) was associated with reduced FL (mean 0.15 z score) and humerus length (mean 0.18 z score) compared to higher exposure (i.e. >50 nmol/L). The associated reduction was limited

to those whose mothers had both reduced plasma 25(OH)D and reduced dietary calcium intake (i.e. <1050 mg/d)(Young BE et al. 2012); this study was at high risk for recruitment bias and probably low risk for bias in two other domains of bias. Two studies from the SWS(Mahon P et al. 2010; Ioannou C et al. 2012), one with probably low risk for bias in confounding and probably high risk for bias due to incomplete outcome(Mahon P et al. 2010), used 3-D ultrasound technology to explore associations between maternal plasma vitamin D status at 34 weeks gestation and fetal femur dimensions. There was no association between maternal vitamin D and FL but there were associations with characteristics of the femur. One study, which included data from 424 fetuses, found an association between reduced maternal 25(OH)D (i.e. <25 nmol/L) and greater splaying of the distal femur at 19 and 34 weeks(Mahon P et al. 2010). The second study(Ioannou C et al. 2012) reported a weak positive association between maternal 25 (OH)D and proximal metaphyseal diameter ($r=0.18$), but not estimated femur volume at 34 weeks gestation in 357 fetuses.

Vitamin E. Maternal plasma α -tocopherol (vitamin E) was positively linked to CRL at ten weeks gestation in 766 mother-fetus pairs: the mean CRL was 42mm and 46 mm for the lowest and highest quartiles, respectively(Turner SW et al. 2010). Maternal exposure was not linked to fetal size later in gestation. The study was at probably low risk for bias in incomplete outcome.

Fatty Acids. A Danish study related maternal whole blood fatty acid composition at 24 weeks gestation to fetal measurements made at 20 weeks gestation in 583 fetuses(Carlsen K et al. 2013), and found a weak relationship between increasing n-3 PUFA and reducing FL (but not HC or AC), the regression coefficient was -0.15 $p=0.02$.

Folate, vitamin B12 and homocysteine. A study from the Generation R cohort related maternal and cord plasma concentrations of folate and vitamin B12, and also homocysteine (associated with reduced birth weight and antioxidant properties) at 13 weeks gestation to fetal weight gain and size at birth(Bergen et al. 2016). The main finding was that lower folate and higher homocysteine in maternal and cord plasma were associated with a slowing of

fetal growth as pregnancy progressed (equivalent to 0.3 z score reduction in birth weight between the upper and lower quartile groups). Unexpectedly, the highest quintile of maternal and cord plasma vitamin B12 was associated with lower birth weight compared to the lowest quintile (mean difference 43g for maternal plasma and 258g for cord plasma). A second Generation R study (Timmermans S et al. 2009) related timing of folate supplementation to EFW in 6365 fetuses and, consistent with the previously mentioned study (Bergen et al. 2016), found increased second trimester AC, third trimester AC and HC and EFW growth for those whose mothers started folic acid before or at the time of conception relative to no folic acid (mean EFW difference 0.10 z scores [0.02, 0.19]); individuals whose mothers who started folate supplementation in the first eight weeks had greater EFW growth in the second and third trimester (but no other fetal measurements) compared to no folic acid. A third Generation R study linked plasma folate concentration to fetal head growth and identified a small positive association equivalent to 3mm at birth per standard deviation increase in folate concentrations (Steenweg-de Graaff et al. 2017).

Maternal dietary intake and fetal measurements

Energy rich diet. A study from the Generation R cohort (Bouwland-Both MI et al. 2013) reported that an energy rich diet (i.e. rich in bread, nuts and margarine and therefore vitamins D and E), was associated with increased size in first trimester in 847 fetuses (mean increase in CRL 1.6mm relative to low energy rich diet); there was no association between diet and fetal measurements in later pregnancy. An observational study from Pakistan recruited 240 pregnant women (80 in each trimester, half of whom who fasted and half who did not fast during the month of Ramadan) and although mothers who fasted were 1kg lighter in the second and third trimester, there was no difference in growth in FL, BPD or EFW in any of the trimesters (Karateke et al. 2015); this study was at probably high risk for bias for confounding since socioeconomic status was not considered.

Mediterranean diet. A study from the Generation R cohort (Timmermans S et al. 2012) linked maternal Mediterranean diet to fetal measurements (n=3207), and individuals whose

mothers were in the tertile with highest adherence to a “Mediterranean diet” (characterised by higher intake of fruit, vegetables, fish, pasta and rice) had higher EFW at 20 and 30 weeks and increased AC at 30 20 weeks compared to those in the lowest adherence tertile (mean difference in z scores 0.11 for both measurements). The study was at probably high risk for bias in exposure assessment and probably low risk from incomplete data.

Milk. A further Generation R cohort paper described associations between maternal milk intake, as reported by food frequency questionnaire at 13 weeks gestation, and fetal measurements at 21 and 30 weeks gestation in 3405 mother-fetus pairs(Heppe DH et al. 2011b). There were associations between higher intakes and increased second trimester EFW and third trimester HC; with reference to ≤ 1 glass/day, 1-2 glasses/day was associated with 0.8mm increased HC and a 6g increased EFW. The study was at probably high risk for bias in exposure assessment and incomplete data.

Fish and seafood. A further Generation R study found no association between maternal fish and seafood consumption to fetal measurements at 21 and 30 weeks(Heppe DH et al. 2011a). A report from the EDEN mother–child cohort also found no association between maternal seafood intake and fetal measurements (Drouillet P et al. 2009). A second report from this cohort found no consistent evidence for mercury contamination of seafood to be associated with reduced second or third trimester fetal size in 691 mother-child pairs where maternal hair mercury was assessed (Drouillet-Pinard P et al. 2010). Both reports from the EDEN study were scored probably low risk for incomplete outcome since they reported significant associations between exposure and outcome for the subset whose mothers were overweight; this analysis was not pre-specified and the findings may be false positives.

Quality and strength of evidence

Collectively the quality of evidence linking maternal dietary exposures to fetal measurements was low since there was low risk of bias in population recruitment in all but one population, and often further risk from (questionnaire-based) exposure assessment and from incomplete outcomes. The strength of evidence was inadequate to relate exposure to fetal size and

growth due to lack of replication for most exposures and differences in methodology between studies which did consider the same dietary exposure.

Ambient air pollution

Ten studies described associations between maternal air pollution exposure and fetal ultrasound scan measurement. Three studies were from the INMA cohort. Air pollutant exposures were: nitrogen dioxide (NO₂), particulate matter <10 microns (PM₁₀), benzene, sulphur dioxide (SO₂), carbon monoxide (CO) and ozone (O₃). Table 2 provides further details of the methodology used and magnitude of any association.

Nitrogen dioxide. Nine studies described the relationship between maternal NO₂ exposure and fetal measurement. Seven studies described associations between higher NO₂ exposures and reduced fetal size and/or growth(Aguilera I et al. 2010; Clemens et al. 2017; Iniguez C et al. 2012; Iniguez et al. 2016; Malmqvist et al. 2017; Ritz B et al. 2014; van den Hooven EH et al. 2012) and two studies reported no association(Carvalho M.A. et al. 2016; Hansen CA et al. 2008). The first of three studies from the INMA cohort, limited to 562 mothers in one of the recruitment centres (Sabadell), found increased NO₂ exposure was associated with a restriction in HC growth at weeks 12-20 and a reduction in BPD, intAC and EFW restriction at weeks 20-32 but only among mothers who spent <2 hours outdoors in non-residential areas per day(Aguilera I et al. 2010). The second report from the INMA cohort(Iniguez C et al. 2012) extended their previous results(Aguilera I et al. 2010) by relating cumulative NO₂ exposure during pregnancy to fetal size in a larger proportion of the cohort (785 mother). Here, the authors demonstrated an association for all mothers between increased maternal NO₂ exposure (i.e. 38 µg/m³) and a reduction in growth for BPD, AC and EFW between 20 and 34 weeks gestation; the authors concluded that NO₂ exposure before 20 weeks gestation was critical to the associations described. The third paper from the INMA cohort extended the previous findings(Iniguez C et al. 2012) to 2478 mothers(Iniguez et al. 2016) and the latter found evidence of faltering growth in the second

(as well as the third) trimester associated with increasing NO₂ exposure. Two papers observed associations between increasing maternal NO₂ exposure and reduced fetal head size, but no other fetal measurement(Clemens et al. 2017; Ritz B et al. 2014). One study reported increased maternal NO₂ exposure was associated with both reduced fetal head size and FL(van den Hooven EH et al. 2012) and another with both reduced FL and AC but not HC or BPD at 32-33 weeks gestation(Malmqvist et al. 2017). A small study of 366 mothers where NO₂ exposure was measured using personal passive samplers in each trimester found no association between exposure and third trimester EFW(Carvalho M.A. et al. 2016).

Fine particulates (PM₁₀). Three of the four papers which related maternal PM₁₀ exposure to fetal measurements found a link between increasing exposure and smaller measurements(Clemens et al. 2017; Hansen CA et al. 2008; van den Hooven EH et al. 2012). Where associations were present, they were found with reduced head size. The association with reduced head size was present only in the second trimester in one study(Hansen CA et al. 2008) and only after the second trimester in the other two studies(Clemens et al. 2017; van den Hooven EH et al. 2012). The study by Clemens *et al*(Clemens et al. 2017) found the association with PM₁₀ was also present for smaller particulates (PM_{2.5}) and also that the association was restricted to fetuses of non-smoking mothers. One study found an association between PM₁₀ exposure and reduced FL(Hansen CA et al. 2008) and a fourth found no association with any fetal measurement(Ritz B et al. 2014). The study by Hansen *et al* (Hansen CA et al. 2008) did not include socioeconomic status was rated probably high risk for bias in confounding.

Benzene. Two studies described an association between increased maternal benzene exposure and reduced head size(Aguilera I et al. 2010; Slama R et al. 2009). One study which was restricted to non-smoking mothers reported reduced second and third trimester fetal head size associated with increasing benzene exposure (equivalent to 2mm difference by 35 weeks gestation)(Slama R et al. 2009). The second study found no association

between benzene and other fetal measurements including FL, AC and EFW, but found reduced BPD growth between weeks 20-32 associated with benzene exposure among a subset of mothers who did not spend ≥ 15 hours at home per day (Aguilera I et al. 2010).

Other exposures. One study found an association between maternal ozone exposure in the first trimester and AC (Hansen CA et al. 2008) but two other studies found no association between ozone exposure and any fetal measurement (Carvalho M.A. et al. 2016; Ritz B et al. 2014). Only one study related SO_2 exposure to fetal measurements and reported associations between increased first trimester exposure and reduced BPD and AC (Hansen CA et al. 2008). One study explored the relationship between CO and fetal measurements and found no associations (Ritz B et al. 2014).

Quality and strength of evidence

Collectively the ten well-designed well-conducted studies provided a high quality of evidence linking maternal exposure to NO_2 and $\text{PM}_{2.5}$ to reduced third trimester fetal measurements (especially head size). The “default” for human studies having moderate quality of evidence was upgraded on account of evidence of dose response between NO_2 and $\text{PM}_{2.5}$ exposures and reduced fetal head size. Many studies were at probably low risk for bias in recruitment and exposure assessment. There was sufficient evidence to link NO_2 and $\text{PM}_{2.5}$ to reduced fetal head size since findings were consistent across a number of studies, reporting different concentrations of exposures, using different design and in different countries. The evidence linking other air pollution exposures to fetal measurements was of high quality but there was inadequate evidence of an association due to inconsistent findings between studies (e.g. benzene) and not all exposures were measured in all studies, e.g. ozone, carbon monoxide.

Maternal alcohol intake

One paper from the Generation R cohort assessed maternal alcohol intake in the first, second and third trimesters for 7333 mother-fetal pairs (Bakker R et al. 2010). In the

longitudinal analysis, the fetuses of mothers who continued to consume alcohol (typically <1 drink a week) had a small increase in EFW growth (0.6g per week). In cross-sectional analyses, there was no association between second and third trimester intake and fetal HC, AC, FL or EFW. There was no evidence of a dose-response effect of alcohol on fetal measurements. A second publication which excluded mothers who smoked obtained routinely collected ultrasound measurements between 18 and 41 weeks gestation and reported reduced HC:AC during the pregnancy for fetuses whose mothers who continued to drink compared to those who quit (magnitude not described)(Handmaker NS et al. 2006). There was reduced cerebellar growth, but no difference in HC, AC and FL, among fetuses exposed to persistent maternal alcohol intake compared to those whose mothers who quit drinking during pregnancy. In this cohort(Handmaker NS et al. 2006), 40% of mothers were marijuana users and 20% used amphetamines with the latter being associated with increased HC:AC growth. A third study (a pilot study) screened 6745 Ukrainian mothers in early pregnancy and identified 84 moderate-to-heavy drinkers and 82 abstinent mothers(Kfir M et al. 2009); moderate-to-heavy drinking was a reported average of 30mls ethanol (or three UK “units”) daily at conception and 4mls daily after knowingly being pregnant (including binges with an average of 66mls ethanol/day). FL was reduced in the second trimester among alcohol exposed fetuses (mean 53rd centile versus 65th centile for controls) but not in the third trimester. Third trimester BPD was reduced in exposed fetuses when compared to controls (54th centile versus 70th centile, respectively).

Collectively these studies were of low quality due to bias in recruitment, use of reported alcohol intake and incomplete follow up and within this limited literature there was inadequate evidence identified to link maternal alcohol intake to fetal measurements.

Other exposures

Occupational exposures. A study of occupational exposures (by questionnaire) in 4680 pregnant mothers related FL, HC and EFW to exposures including phthalates, pesticides,

polyaromatic hydrocarbons and alkylphenolic compounds(Snijder CA et al. 2012b). There were associations of small magnitude seen between some exposures and some fetal measurements, most consistently for phthalates whose exposure was associated with reduced EFW and fetal length (approximately 1% of a z score per week).

Bisphenol A. A second Generation R study measured maternal urinary Bisphenol A (BPA) in the first, second and third trimesters and related these to fetal growth in 419 pregnancies (Snijder CA et al. 2013). In 80 mothers (i.e. 19% of the study population) urine samples were obtained in each trimester there was reduced growth EFW and HC per unit increase in BPA (typically 2% reduction per week per unit increase in BPA exposure), but this finding was not replicated in models which included mothers where exposure data were not complete nor in a report from the INMA cohorts where there was no association between maternal first and third trimester BPA and fetal measurements (488 mothers)(Casas et al. 2016).

Phenols. A study from the EDEN cohort related maternal exposure to nine phenol-based chemicals to fetal size and growth throughout pregnancy and found evidence that one (triclosan) was linked to minor reductions in fetal size in the third trimester but at no other time(Philippat et al. 2014).

Phthalates. Two studies related maternal phthalate exposure to fetal size and growth and report apparently contrasting findings. The first study, again restricted to male offspring in the EDEN cohort, measured 11 phthalate molecules in maternal urine and found a negative relationship between phthalate metabolites and EFW (10-15g per quartile increase in phthalate exposure) but also a positive association between one molecule (monocarboxyisononyl) and FL in the second and third trimesters(Botton et al. 2016) (which is consistent with the previously mentioned study(Casas et al. 2016)). A study from the INMA cohorts found association of small magnitude between increased concentrations of two of eight phthalate molecules in maternal urine and reduced growth HC in the first, and FL second half of pregnancy(Casas et al. 2016). Both cohorts only measured one common phthalate molecule (monoethyl phthalate) so the results are not necessarily inconsistent.

Polybrominated diphenyl ethers (PBDEs). A further evaluation of the INMA cohorts measured PBDEs in maternal and cord concentrations and related these to fetal measurements in 670 mothers (Lopez-Espinosa et al. 2015); a doubling of PBDE exposure was associated with <5% reduced growth in AC, EFW and BPD (but not FL) between 20-34 weeks (but not 12-20 weeks), although associations were not always present for both maternal and cord blood concentrations.

Organochlorine compounds. The final study identified from 2369 the INMA cohorts related maternal organochlorine compounds in maternal plasma at 12 weeks gestation and cord blood to fetal growth in early, mid and late pregnancy (Lopez-Espinosa et al. 2016); a doubling of exposures to polychlorinated biphenyls (PCB) -138, -153 and -180 were associated with a 2-4% reduction in femur length growth and a doubling of PCB-138 was associated with a 2% reduction in EFW growth between gestational weeks 20-34. There were inverse associations between cord Hexachlorobenzene and reduced AC growth between 0 and 20 weeks. Five of 120 associations in this analysis achieved significance.

Lithium in drinking water. The northern Argentinian Andes has variable concentrations of lithium in its drinking water and a study of 194 mothers observed no significant relationships between maternal plasma and urinary lithium and second and third trimester BPD, HC, AC, FL and EFW (Harari et al. 2015).

Collectively the quality of evidence for these studies was low since although objective measurements of exposure were made in all but one study, multiple comparisons were made within studies increasing the chance of false positive findings, findings (when replicated) were not consistent and risk of bias from incomplete follow up was probably high or high in four studies. Additionally, data were limited only to male offspring in two studies (Botton et al. 2016; Philippat et al. 2014). In conclusion, there was inadequate strength of evidence linking chemicals to fetal size since the quality of evidence was low, many different chemicals were studied and where associations with fetal size were present, these were not replicated elsewhere.

DISCUSSION

This systematic review was designed to describe the literature associating maternal exposures to fetal size being small for gestational age. The papers identified were all published since 2006, indicating that this is a relatively new literature. The first major finding was that maternal exposures to increased ambient air NO₂ and PM_{2.5} were consistently associated with reduced third trimester fetal head size and we judge that the literature presents sufficient evidence of toxicity in this context. The second major finding was that the strength of evidence was inadequate for all the other exposures considered. The third notable finding was that, where associations were present, they were more commonly seen in the third compared to the second trimester and that reduced third trimester head size was more commonly associated with potentially adverse maternal exposures compared to other fetal measurements. Together these findings suggest that public health measures are urgently required to minimise pregnant mother's exposure to NO₂ and PM_{2.5}, and more high-quality research is required to better understand the relationship between other (modifiable) maternal exposures and fetal measurements.

The fetus has traditionally been thought to have a privileged position, where it was protected from the adverse effects of environmental exposures by the maternal-placental "unit", but associations between reduced birth weight and many maternal exposures argue against this paradigm. The fetal origins hypothesis (Barker 1995a) speculated that maternal exposures in mid pregnancy were relevant to birth weight and risk for subsequent increased risk for non-communicable diseases. Since we find evidence of reduced second trimester fetal size and some maternal exposures, and our findings suggest that maternal exposures in early pregnancy are also relevant.

There were some associations between exposure and fetal size which were mostly consistent across different populations, whilst other associations were less consistently seen, or were even counterintuitive. An example of consistency was the inverse association

between NO₂ exposure and third trimester fetal size, which was seen in all seven studies which explored this link (two studies measured NO₂ but not third trimester head size Carvalho M.A. et al. 2016; Hansen CA et al. 2008). In contrast, there were some instances where an apparently harmful exposure was associated with increased fetal size (e.g. current PM₁₀ exposure and first trimester length and second trimester weight (van den Hooven EH et al. 2012), one phthalate molecule and increased third trimester fetal length (Botton et al. 2016), increased maternal and cord plasma vitamin B12 concentrations and reduced birth weight (Bergen et al. 2016), maternal alcohol intake and increased third trimester EFW (Bakker R et al. 2010)); although these associations may challenge the paradigm that “harmful” exposures invariably cause reduced fetal growth, these may be false positive findings.

A further example of inconsistency comes from the ambient air exposure literature where two studies found that the magnitude of association between NO₂ and fetal size was only present (Ritz B et al. 2014), or was greater (Iniguez et al. 2016), among mothers who smoked whereas a second study only found an association between NO₂ and fetal head size among non-smokers (Clemens et al. 2017). Our review also identified many associations of borderline significance and instances of multiple testing, and these increase the risk of false positive results. Given that the body of literature reviewed is relatively young, it is not unexpected that there is a spectrum of consistency/inconsistency between maternal exposure and fetal size and further research activity is required to replicate some of the apparently inconsistent associations.

There are consistencies and inconsistencies in the literature describing associations between maternal exposures and birth weight and these exposures and fetal size. Maternal exposures to poor quality ambient air is associated with reduced birth weight (Maisonet et al. 2004; Shah PS. Balkhair T. Knowledge Synthesis Group on Determinants of Preterm/LBW births 2011) and we confirm this association is already present in fetal life. In contrast, many maternal dietary exposures are associated with reduced birth weight (Gresham E et al.

2014) but there were no consistent associations between maternal dietary exposures and fetal size in the literature we reviewed. Similarly, maternal exposure to chemicals such as bisphenol A(Perera F and Herbstman J 2011) is associated with reduced birth weight but bisphenol A was associated with reduced fetal size in one study (Casas et al. 2016) but not a second (Snijder CA et al. 2013). The inconsistent findings between size before and after birth in the context of maternal exposures may reflect a smaller literature describing antenatal size, the challenges of accurately measuring fetal size or theoretically may be due to associations only becoming detectable towards the very end of pregnancy.

Maternal alcohol ingestion during pregnancy is an established risk factors for reduced birth weight(Henderson J et al. 2007), but the literature was not adequate to determine when the relationship between alcohol ingestion and fetal growth failure begins. The studies identified were at high risk of bias and collectively provided inadequate evidence of toxicity This absence of evidence does not mean that any change is required to current guidelines which recommend that pregnant mothers should abstain from drinking alcohol.

A weakness of the present literature is that all the evidence comes from observational studies where, in addition to the risk of false positive findings previously discussed, some of the results may be influenced by confounding factors and also the results are based on populations who are not necessarily representative of the general population. Drop out from cohort studies also may contribute bias, for example a study which related fetal size to risk for later asthma found a relationship of greater magnitude when using questionnaire reported asthma outcome (available in 39% of the cohort) compared to using routinely acquired asthma outcome (available in 88%) (Turner et al. 2018).

One further limitation of the literature is that whilst there are some maternal exposures linked to fetal size in three or more studies, for example smoking (Abraham et al. 2017) and air pollution (the present review), many exposures are linked to fetal size in only one or two studies (for example chemicals and dietary nutrients) and this leaves insufficient evidence

upon which to form an opinion. A second potential limitation to the literature is that fetal measurements and exposures were made at different gestational age, although this limitation applies to the different studies in the literature linking post natal exposures and post natal outcomes. Fetal size is driven by gestational age, and fetal size was determined over a range of gestations within and between different studies. Gestational age was included as a covariate in all but one study (Karateke et al. 2015) and this minimises the potential for differences in gestational age at assessment confounding associations with maternal exposure. Differences between studies in the gestational age when fetal size and maternal exposure are likely to weaken and not strengthen the associations (where present) between maternal exposure and fetal size.

AOne further limitation to the literature was that different methods were used to measure exposures, even where the same exposure (NO₂ for example) was measured in several studies. This meant that meta-analysis was not possible for single exposures. ~~A second limitation of the literature is that many of the exposures studied, especially maternal diet and plasma nutrients, there was only one study which linked that exposure to fetal size and the findings require replication.~~ Another limitation to the literature is that there are some areas where the evidence is of a poor quality, e.g. maternal alcohol exposure, which is an increasing public health concern due to increasing awareness of fetal alcohol syndrome (Popova et al. 2017). Finally, there are several maternal exposures including recreational drug use (Behnke M et al. 2013), pesticides (Sathyanarayana et al. 2010), proximity to land fill sites (Elliott et al. 2001) and electromagnetic fields (de Vocht F and Lee B 2014) which are linked to reduced birth weight but where there are no studies associating such exposures to fetal measurements.

Pregnant mothers are constantly exposed to multiple environmental exposures and identifying which single exposure may be important to fetal size is a challenge, for example increased maternal alcohol intake may be associated with smoking, use of illicit drugs and poor diet. Furthermore, any reduction in fetal size associated with tobacco and alcohol

consumption may be modified by increased size associated with amphetamine intake (Handmaker NS et al. 2006). Only intervention studies are able to infer causality, but these are practically and ethically challenging to do in the context of potentially harmful maternal exposures during pregnancy and the evidence base may remain dependent on observational studies.

The mechanism(s) where environmental exposures may cause fetal growth failure (or acceleration) are not well understood but likely to be multiple. The mechanism for maternal diet influencing fetal size assumes that macro and micro nutrients cross the placenta, and maternal dietary fat intake during the third trimester has been associated with cord blood lipid concentrations (Elias SL and Innis SM 2001) but the fetus is not necessarily exposed to all maternal plasma nutrients, for example triglycerides do not cross the placenta (Herrera 2002). The mechanism for ambient air exposures leading to fetal growth failure is not understood but ultrafine particles (which correlate with PM₁₀ and NO₂ concentrations) are able to directly enter the maternal circulation via the alveoli (Geiser M et al. 2005) and placenta (Wick P et al. 2010), and thus theoretically enter the fetal circulation where they may be pathogenic by causing vascular inflammation (Gojova A et al. 2007).

In summary, there is an emerging literature which suggests that some maternal exposures during early pregnancy are associated with changes in fetal size which are apparent by the second and third trimester. Evidence from randomised controlled trials where maternal exposures are modified are unlikely due to practical and ethical issues, and the evidence base is likely to arise from observational studies. The literature is mostly based on Western populations and is dominated by three cohort studies and replication in other populations of the associations described is required.

ACKNOWLEDGEMENTS

The authors thank Dr Juleen Lam for her advice on the use of the Navigation guide and for her helpful comments on a draft of the submitted manuscript.

REFERENCES

- Abraham, M.; Alramadhan, S.; Iniguez, C.; Duijts, L.; Jaddoe, V.W.V.; Den Dekker, H.T. et al. A systematic review of maternal smoking during pregnancy and fetal measurements with meta-analysis. *PLoS ONE* 12:e0170946; 2017.
- Aguilera I; Garcia-Esteban R; Iniguez C; Nieuwenhuijsen MJ; Rodriguez A; Paez M et al. Prenatal exposure to traffic-related air pollution and ultrasound measures of fetal growth in the INMA Sabadell cohort. *Environ Health Perspect* 118:705-11; 2010.
- Alkandari, F.; Ellahi, A.; Aucott, L.; Devereux, G.; Turner, S. Fetal ultrasound measurements and associations with postnatal outcomes in infancy and childhood: a systematic review of an emerging literature. *J Epidemiol Comm Health* 69:41-48; 2015.
- Bakker R; Pluimgraaff LE; Steegers EA; Raat H; Tiemeier H; Hofman A et al. Associations of light and moderate maternal alcohol consumption with fetal growth characteristics in different periods of pregnancy: the Generation R Study. *Int J Epidemiol* 39:777-89; 2010.
- Barker, D.J. Fetal origins of coronary heart disease. *BMJ* 311:171-4; 1995.
- Bateson, P.; Barker, D.; Clutton-Brock, T.; Deb, D.; D'Udine, B.; Foley, R.A. et al. Developmental plasticity and human health. *Nature* 430:419-21; 2004.
- Behnke M; Smith VC; Committee on Substance Abuse; Committee on Fetus and Newborn. Prenatal substance abuse: short- and long-term effects on the exposed fetus. *Pediatrics* 131:e1009-24; 2013.
- Bergen, N.E.; Schalekamp-Timmermans, S.; Jaddoe, V.W.V.; Hofman, A.; Lindemans, J.; Russcher, H. et al. Maternal and Neonatal Markers of the Homocysteine Pathway and Fetal Growth: The Generation R Study. *Paediatr Perinat Epidemiol* 30:386-96; 2016.
- Botton, J.; Philippat, C.; Calafat, A.M.; Carles, S.; Charles, M.; Slama, R. et al. Phthalate pregnancy exposure and male offspring growth from the intra-uterine period to five years of age. *Environ Res* 151:601-9; 2016.
- Bouwland-Both MI; Steegers-Theunissen RP; Vujkovic M; Lesaffre EM; Mook-Kanamori DO; Hofman A et al. A periconceptional energy-rich dietary pattern is associated with early fetal growth: the Generation R study. *BJOG: An International Journal of Obstetrics & Gynaecology* 120:435-45; 2013.
- Carlsen K; Pedersen L; Bonnelykke K; Stark KD; Lauritzen L; Bisgaard H. Association between whole-blood polyunsaturated fatty acids in pregnant women and early fetal weight. *Eur J Clin Nutr* 67:978-83; 2013.
- Carvalho M.A.; Bernardes L.S.; Hettfleisch K.; Pastro L.D.M.; Vieira S.E.; Saldiva S.R.D.M. et al. Associations of maternal personal exposure to air pollution on fetal weight and fetoplacental Doppler: A prospective cohort study. *Reproductive Toxicology* 62:9-17; 2016.
- Casas, M.; Valvi, D.; Ballesteros-Gomez, A.; Gascon, M.; Fernandez, M.F.; Garcia-Esteban, R. et al. Exposure to Bisphenol A and Phthalates during Pregnancy and Ultrasound

- Measures of Fetal Growth in the INMA-Sabadell Cohort. *Environ Health Perspect* 124:521-8; 2016.
- Clemens, T.; Turner, S.; Dibben, C. Maternal exposure to ambient air pollution and fetal growth in North-East Scotland: A population-based study using routine ultrasound scans *Environ Int* 107:216-226; 2017.
- de Vocht F and Lee B. Residential proximity to electromagnetic field sources and birth weight: Minimizing residual confounding using multiple imputation and propensity score matching. *Environ Int* 69:51-7; 2014.
- Drouillet P; Kaminski M; De Lauzon-Guillain B; Forhan A; Ducimetiere P; Schweitzer M et al. Association between maternal seafood consumption before pregnancy and fetal growth: evidence for an association in overweight women. The EDEN mother-child cohort. *Paediatr Perinat Epidemiol* 23:76-86; 2009.
- Drouillet-Pinard P; Huel G; Slama R; Forhan A; Sahuquillo J; Goua V et al. Prenatal mercury contamination: relationship with maternal seafood consumption during pregnancy and fetal growth in the 'EDEN mother-child' cohort. *Br J Nutr* 104:1096-100; 2010.
- Elias SL and Innis SM. Infant plasma trans, n-6, and n-3 fatty acids and conjugated linoleic acids are related to maternal plasma fatty acids, length of gestation, and birth weight and length. *Am J Clin Nutr* 73:807-14; 2001.
- Elliott, P.; Briggs, D.; Morris, S.; de Hoogh, C.; Hurt, C.; Jensen, T.K. et al. Risk of adverse birth outcomes in populations living near landfill sites. *BMJ* 323:363-8; 2001.
- Geiser M; Rothen-Rutishauser B; Kapp N; Schurch S; Kreyling W; Schulz H et al. Ultrafine particles cross cellular membranes by nonphagocytic mechanisms in lungs and in cultured cells. *Environ Health Perspect* 113:1555-60; 2005.
- Gluckman, P.D.; Hanson, M.A.; Low, F.M. The role of developmental plasticity and epigenetics in human health. *Birth Defects Research. Part C, Embryo Today: Reviews* 93:12-8; 2011.
- Gluckman, P.D.; Hanson, M.; Spencer, H.G. Predictive adaptive responses and human evolution. *Trends Ecol Evol* 20:527-33; 2005.
- Gojova A; Guo B; Kota RS; Rutledge JC; Kennedy IM; Barakat AI. Induction of inflammation in vascular endothelial cells by metal oxide nanoparticles: effect of particle composition. *Environ Health Perspect* 115:403-9; 2007.
- Gresham E; Byles JE; Bisquera A; Hure AJ. Effects of dietary interventions on neonatal and infant outcomes: a systematic review and meta-analysis. *Am J Clin Nutr* 100:1298-321; 2014.
- Guxens M; Ballester F; Espada M; Fernandez MF; Grimalt JO; Ibarluzea J et al. Cohort Profile: the INMA--Infancia y Medio Ambiente--(Environment and Childhood) Project. *Int J Epidemiol* 41:930-40; 2012.
- Hadlock, F.P.; Harrist, R.B.; Carpenter, R.J.; Deter, R.L.; Park, S.K. Sonographic estimation of fetal weight. The value of femur length in addition to head and abdomen measurements. *Radiology* 150:535-40; 1984.

- Hales CN and Barker DJ. Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis. *Diabetologia* 35:595-601; 1992.
- Handmaker NS; Rayburn WF; Meng C; Bell JB; Rayburn BB; Rappaport VJ. Impact of alcohol exposure after pregnancy recognition on ultrasonographic fetal growth measures. *Alcoholism: Clinical & Experimental Research* 30:892-8; 2006.
- Hansen CA; Barnett AG; Pritchard G. The effect of ambient air pollution during early pregnancy on fetal ultrasonic measurements during mid-pregnancy. *Environ Health Perspect* 116:362-9; 2008.
- Harari, F.; Langeen, M.; Casimiro, E.; Bottai, M.; Palm, B.; Nordqvist, H. et al. Environmental exposure to lithium during pregnancy and fetal size: a longitudinal study in the Argentinean Andes. *Environ Int* 77:48-54; 2015.
- Henderson J; Kesmodel U; Gray R. Systematic review of the fetal effects of prenatal binge-drinking. *Journal of Epidemiology & Community Health* 61:1069-73; 2007.
- Heppe DH; Steegers EA; Timmermans S; Breeijen Hd; Tiemeier H; Hofman A et al. Maternal fish consumption, fetal growth and the risks of neonatal complications: the Generation R Study. *Br J Nutr* 105:938-49; 2011a.
- Heppe DH; van Dam RM; Willemsen SP; den Breeijen H; Raat H; Hofman A et al. Maternal milk consumption, fetal growth, and the risks of neonatal complications: the Generation R Study. *Am J Clin Nutr* 94:501-9; 2011b.
- Herrera, E. Implications of dietary fatty acids during pregnancy on placental, fetal and postnatal development--a review. *Placenta* 23:S9-19; 2002.
- Iniguez C; Ballester F; Estarlich M; Esplugues A; Murcia M; Llop S et al. Prenatal exposure to traffic-related air pollution and fetal growth in a cohort of pregnant women. *Occupational & Environmental Medicine* 69:736-44; 2012.
- Iniguez, C.; Esplugues, A.; Sunyer, J.; Basterrechea, M.; Fernandez-Somoano, A.; Costa, O. et al. Prenatal Exposure to NO₂ and Ultrasound Measures of Fetal Growth in the Spanish INMA Cohort. *Environ Health Perspect* 124:235-42; 2016.
- Ioannou C; Javaid MK; Mahon P; Yaqub MK; Harvey NC; Godfrey KM et al. The effect of maternal vitamin D concentration on fetal bone. *Journal of Clinical Endocrinology & Metabolism* 97:E2070-7; 2012.
- Johnson, P.I.; Sutton, P.; Atchley, D.S.; Koustas, E.; Lam, J.; Sen, S. et al. The Navigation Guide - evidence-based medicine meets environmental health: systematic review of human evidence for PFOA effects on fetal growth. *Environ Health Perspect* 122:1028-39; 2014.
- Karateke, A.; Kaplanoglu, M.; Avci, F.; Kurt, R.K.; Baloglu, A. The effect of Ramadan fasting on fetal development.. *Pakistan Journal of Medical Sciences* 31:1295-9; 2015.
- Kfir M; Yevtushok L; Onishchenko S; Wertelecki W; Bakhireva L; Chambers CD et al. Can prenatal ultrasound detect the effects of in-utero alcohol exposure? A pilot study. *Ultrasound in Obstetrics & Gynecology* 33:683-9; 2009.

Lopez-Espinosa, M.; Costa, O.; Vizcaino, E.; Murcia, M.; Fernandez-Somoano, A.; Iniguez, C. et al. Prenatal Exposure to Polybrominated Flame Retardants and Fetal Growth in the INMA Cohort (Spain). *Environ Sci Technol* 49:10108-16; 2015.

[Lopez-Espinosa, M.; Murcia, M.; Iniguez, C.; Vizcaino, E.; Costa, O.; Fernandez-Somoano, A. et al. Organochlorine Compounds and Ultrasound Measurements of Fetal Growth in the INMA Cohort \(Spain\). *Environ Health Perspect* 124:157-163; 2016](#)

Mahon P; Harvey N; Crozier S; Inskip H; Robinson S; Arden N et al. Low maternal vitamin D status and fetal bone development: cohort study. *Journal of Bone & Mineral Research* 25:14-9; 2010.

Maisonet, M.; Correa, A.; Misra, D.; Jaakkola, J.J. A review of the literature on the effects of ambient air pollution on fetal growth. *Environ Res* 95:106-15; 2004.

Malmqvist, E.; Liew, Z.; Kallen, K.; Rignell-Hydbom, A.; Rittner, R.; Rylander, L. et al. Fetal growth and air pollution - A study on ultrasound and birth measures. *Environ Res* 152:73-80; 2017.

Pereira, P.P.d.S.; Da Mata, F.A.F.; Figueiredo, A.C.G.; de Andrade, K.R.C.; Pereira, M.G. Maternal Active Smoking During Pregnancy and Low Birth Weight in the Americas: A Systematic Review and Meta-analysis. *Nicotine Tobacco Res* 19:497-505; 2017.

Perera F and Herbstman J. Prenatal environmental exposures, epigenetics, and disease. *Reproductive Toxicology* 31:363-73; 2011.

Philippat, C.; Botton, J.; Calafat, A.M.; Ye, X.; Charles, M.; Slama, R. et al. Prenatal exposure to phenols and growth in boys.. *Epidemiology* 25:625-35; 2014.

Popova, S.; Lange, S.; Probst, C.; Gmel, G.; Rehm, J. Estimation of national, regional, and global prevalence of alcohol use during pregnancy and fetal alcohol syndrome: a systematic review and meta-analysis.. *The Lancet Global Health* 5:e290-9; 2017.

Ritz B; Qiu J; Lee PC; Lurmann F; Penfold B; Erin Weiss R et al. Prenatal air pollution exposure and ultrasound measures of fetal growth in Los Angeles, California. *Environ Res* 130:7-13; 2014.

Sathyanarayana, S.; Basso, O.; Karr, C.J.; Lozano, P.; Alavanja, M.; Sandler, D.P. et al. Maternal pesticide use and birth weight in the agricultural health study. *J Agromed* 15:127-36; 2010.

Shah PS. Balkhair T. Knowledge Synthesis Group on Determinants of Preterm/LBW births. Air pollution and birth outcomes: a systematic review. *Environ Int* 37:498-516; 2011.

Shaheen SO; Sterne JA; Montgomery SM; Azima H. Birth weight, body mass index and asthma in young adults. *Thorax* 54:396-402; 1999.

Slama R; Thiebaugeorges O; Goua V; Aussel L; Sacco P; Bohet A et al. Maternal personal exposure to airborne benzene and intrauterine growth. *Environ Health Perspect* 117:1313-21; 2009.

- Snijder CA; Brand T; Jaddoe V; Hofman A; Mackenbach JP; Steegers EA et al. Physically demanding work, fetal growth and the risk of adverse birth outcomes. The Generation R Study. *Occupational & Environmental Medicine* 69:543-50; 2012a.
- Snijder CA; Heederik D; Pierik FH; Hofman A; Jaddoe VW; Koch HM et al. Fetal growth and prenatal exposure to bisphenol A: the generation R study. *Environ Health Perspect* 121:393-8; 2013.
- Snijder CA; Roeleveld N; Te Velde E; Steegers EA; Raat H; Hofman A et al. Occupational exposure to chemicals and fetal growth: the Generation R Study. *Human Reproduction* 27:910-20; 2012b.
- Steenweg-de Graaff, J.; Roza, S.J.; Walstra, A.N.; El Marroun, H.; Steegers, E.A.P.; Jaddoe, V.W.V. et al. Associations of maternal folic acid supplementation and folate concentrations during pregnancy with foetal and child head growth: the Generation R Study.. *Eur J Nutr* 56:65-75; 2017.
- Timmermans S; Jaddoe VW; Hofman A; Steegers-Theunissen RP; Steegers EA. Periconception folic acid supplementation, fetal growth and the risks of low birth weight and preterm birth: the Generation R Study. *Br J Nutr* 102:777-85; 2009.
- Timmermans S; Steegers-Theunissen RP; Vujkovic M; den Breeijen H; Russcher H; Lindemans J et al. The Mediterranean diet and fetal size parameters: the Generation R Study. *Br J Nutr* 108:1399-409; 2012.
- Turner SW; Campbell D; Smith N; Craig LC; McNeill G; Forbes SH et al. Associations between fetal size, maternal {alpha}-tocopherol and childhood asthma. *Thorax* 65:391-7; 2010.
- Turner, S.; Fielding, S.; Devereux, G. First trimester fetal size and prescribed asthma medication at 15 years of age. *European Respiratory Journal* 51; 2018.
- van den Hooven EH; Pierik FH; de Kluizenaar Y; Willemsen SP; Hofman A; van Ratingen SW et al. Air pollution exposure during pregnancy, ultrasound measures of fetal growth, and adverse birth outcomes: a prospective cohort study. *Environ Health Perspect* 120:150-6; 2012.
- Wick P; Malek A; Manser P; Meili D; Maeder-Althaus X; Diener L et al. Barrier capacity of human placenta for nanosized materials. *Environ Health Perspect* 118:432-6; 2010.
- Young BE; McNanley TJ; Cooper EM; McIntyre AW; Witter F; Harris ZL et al. Maternal vitamin D status and calcium intake interact to affect fetal skeletal growth in utero in pregnant adolescents. *Am J Clin Nutr* 95:1103-12; 2012.

FIGURE LEGENDS

Figure 1. CONSORT style diagram showing how the papers included in the review were identified.

Figure two. Bar chart summarising the risk of bias for each of the eight domains considered across the 365 papers included in this systematic review.

Figure three. This figure shows the risk of bias in each of eight domains considered for each of the 365 studies included in this systematic review.

ABSTRACT

Background. Reduced birth weight is associated with many maternal environmental exposures during pregnancy, but the gestational age at onset of this association is unknown.

We have previously reported associations between maternal smoking and fetal size.

Objective. To report on our systematic review of the literature describing associations between antenatal size and growth and maternal exposures during pregnancy.

Data sources. Electronic databases (OVID and EMBASE) and web sites for cohort studies were searched.

Study eligibility. Studies were eligible if they examined associations between maternal environmental exposures (including ambient air exposure, diet and alcohol) and antenatal fetal ultrasound measurements.

Study appraisal. The Navigation Guide was used to assess the strength of evidence.

Results. There were 451 abstracts identified and 36 papers were included of which maternal diet was the exposure of interest in 15, maternal ambient air exposure in 10, maternal alcohol in 3 and other exposures in 8. The first paper was published in 2006. Associations were present between exposures in 18% of comparisons with second trimester measurements and in 46% of comparisons with third trimester measurements. In the third trimester, when an association was present, reduced head size was most commonly (58%) associated with current or previous maternal exposure, with reduced length being least commonly (32%) associated and reduced weight being intermediate (52%). In the third trimester, increased maternal nitrogen dioxide exposure was associated with reduced head size was associated with in all seven studies identified and reduced fetal weight in five out of six studies.

Conclusion. There is sufficient evidence of toxicity in the context of maternal exposure to nitrogen dioxide and reduced third trimester fetal head size. There is insufficient evidence of toxicity with regard to maternal exposures to dietary factors, alcohol and environmental chemicals and reduced fetal size.

Key words. Air pollution; Benzene; Diet; Ethanol; Fetus; Mother; Phthalic acids

ABBREVIATIONS

HC=head circumference

AC=abdominal circumference

BPD=biparietal diameter

BTEX= aromatic hydrocarbons (benzene, toluene, ethylbenzene, m/p-xylene, and o-xylene)

CO=carbon monoxide

4,4'DDE=4,4'dichlorodiphenylchloroethylene

EFW=estimated fetal weight.

HCB= hexachlorobenzene

MAD=mean abdominal circumference

NCD=non-communicable diseases

NO₂= nitrogen dioxide

O₃=ozone

PAH=polycyclic aromatic hydrocarbons

PCB=polychlorinated biphenyls

PM₁₀=particulates with diameter less than 10 microns

PUFA=poly unsaturated fatty acids

SO₂=sulphur dioxide

INTRODUCTION

Being small for gestational age (SGA) at birth is associated with increased risk for conditions that include coronary artery disease(Barker 1995a), type II diabetes(Hales CN and Barker DJ 1992) and asthma(Shaheen SO et al. 1999). The fetal origins hypothesis(Barker 1995a) and more recently, concepts including developmental plasticity(Bateson et al. 2004) and predictive adaptive responses(Gluckman et al. 2005), suggest that some antenatal exposures (or cues) predispose an unborn individual to non-communicable diseases (NCD) in later life by a mechanism which involves reduced fetal size and growth. A recent systematic review linking reduced fetal size, as evidenced by fetal ultrasound scan, to NCD in children supports the principle of the developmental origins of disease(Alkandari et al. 2015).

The mechanism(s) where antenatal cues reduce fetal size and pre-dispose to NCD are not fully understood, but are thought to include epigenetic modification in fetal cells following maternal exposure during pregnancy(Gluckman et al. 2011). Maternal exposures linked to SGA at birth include maternal smoking(Pereira et al. 2017), exposures to poor quality ambient air(Maisonet et al. 2004; Shah PS. Balkhair T. Knowledge Synthesis Group on Determinants of Preterm/LBW births 2011), dietary factors(Gresham E et al. 2014) and exposure to chemicals such as bisphenol A(Perera F and Herbstman J 2011). Knowledge of which exposures are associated with reduced antenatal size and the gestation at which these exposures may be acting would be important to our understanding of “fetal origins” and for public health educational messages and interventions, but the literature describing associations between these exposures and fetal size has not been systematically reviewed. Our group has recently undertaken a systematic review of the literature linking antenatal size and growth to maternal smoking and found that maternal smoking is consistently associated with fetal growth failure after the second trimester(Abraham et al. 2017). Here, we undertake a systematic review designed to answer the population exposure, comparator and outcome (PECO) question: Are fetuses who are exposed to maternal environmental exposures other than smoking small for gestational age compared to unexposed fetuses?

The Navigation Guide methodology is ideally suited for determining the strength of evidence between exposures and fetal size (Johnson et al. 2014) and was used in our study.

METHODS

Search methodology

A database search was carried out September 2017 using OVID MEDLINE and Embase databases and updated in May 2018. Search terms were developed initially using those used in previous systematic reviews (Abraham et al. 2017; Alkandari et al. 2015) and modified after identifying relevant publications not identified using these previous search terms (Aguilera I et al. 2010; Slama R et al. 2009). The online supplement shows the search terms used. Additional papers were identified from bibliographies and from the web sites of the following cohorts known to have collected ultrasound measurements of fetal size: the Raine cohort (<http://www.rainestudy.org.au/>), the EDEN cohort (<https://eden.vjf.inserm.fr>), Southampton Women Study (SWS, <http://www.leu.soton.ac.uk/sws/>), Generation R (<http://www.erasmusmc.nl/epi/research/Generation-R/>), and INMA Mother and Child Cohort Study (Guxens M et al. 2012). The supplement describes these cohorts in greater detail. Two researchers (MA and ST or IH and ST) independently reviewed abstracts identified by the database search and full papers considered potentially eligible were obtained. Eligible papers had to report fetal ultrasound anthropometric measurements (i.e. crown rump length, biparietal diameter, head circumference, femur length, abdominal circumference or estimated fetal weight) and relate these to a current or previous maternal environmental exposures. From background reading, the following exposures were sought: ambient air pollution, maternal alcohol ingestion, maternal diet, maternal drug use, occupational exposures, or pesticide exposure. Papers where fetal size was related to obstetric complications, maternal smoking (as the primary exposure) and maternal drug treatment were excluded. The outcome measures were fetal size and fetal growth.

Fetal measurements. First trimester (i.e. ≤ 13 weeks gestation) measurements were: crown rump length (CRL), biparietal diameter (BPD), head circumference (HC) abdominal

circumference (AC) and mean abdominal diameter (MAD). Second trimester (i.e. >13 to <28 weeks gestation) and third trimester (i.e. ≥28 weeks) measurements were femur length (FL), HC, BPD, MAD, AC and estimated fetal weight (EFW) (the latter was derived from BPD, AC and FL measurements(Hadlock et al. 1984)).

Quality assessment of data

The Navigation Guide methodology was used to assess the risk of bias and the quality and strength of evidence(Johnson et al. 2014). We developed a protocol to assess risk of bias, quality of evidence and strength of evidence and which was based on the study of Johnson *et al* (Johnson et al. 2014) and which is described in the supplement. Risk of bias was rated for each paper and then across each of four exposure groups (i.e. maternal dietary exposures, air pollution exposures, alcohol exposure and other exposures) to inform the rating of quality of the evidence. The quality of the evidence was then used as part of the assessment of the strength of evidence. Two researchers (DM and PC) independently assessed each paper and then consensus was reached during discussion with three researchers (DM, PC and ST).

Meta-Analysis of data

We sought to identify results which might be suitable for meta-analysis. To do this we identified exposures where the same units of exposure were used in studies using comparable methodologies and reporting the same fetal measurement outcome.

RESULTS

Papers identified

The search identified 451 abstracts and 36 papers were ultimately included in this review, (figure 1). Fifteen papers considered maternal dietary factors(Bergen et al. 2016; Bouwland-Both MI et al. 2013; Carlsen K et al. 2013; Drouillet P et al. 2009; Drouillet-Pinard P et al. 2010; Heppe DH et al. 2011a; Heppe DH et al. 2011b; Ioannou C et al. 2012; Karateke et al.

2015; Mahon P et al. 2010; Steenweg-de Graaff et al. 2017; Timmermans S et al. 2009; Timmermans S et al. 2012; Turner SW et al. 2010; Young BE et al. 2012), ten ambient air exposures(Aguilera I et al. 2010; Carvalho M.A. et al. 2016; Clemens et al. 2017; Hansen CA et al. 2008; Iniguez C et al. 2012; Iniguez et al. 2016; Malmqvist et al. 2017; Ritz B et al. 2014; Slama R et al. 2009; van den Hooven EH et al. 2012), three maternal alcohol consumption(Bakker R et al. 2010; Handmaker NS et al. 2006; Kfir M et al. 2009) and eight considered other exposures(Botton et al. 2016; Casas et al. 2016; Harari et al. 2015; Lopez-Espinosa et al. 2015; Philippat et al. 2014; Snijder CA et al. 2012b; Snijder CA et al. 2013; Lopez-Espinosa et al. 2016). Studies which related environmental exposures to fetal anomalies (e.g. kidney cysts) and one relating physical demands of work(Snijder CA et al. 2012a) to fetal size were excluded. All studies were observational and the first was published in 2006. Three publications were from the USA, one each from Australia, Argentina, Brazil, Pakistan and Ukraine and the remaining 28 from European populations. Twenty one publications were from one of three birth cohorts (including ten from the Generation R cohort, six from INMA and five from EDEN).

Risk of bias

There was generally a low risk of overall bias across the studies, with the exception of recruitment bias and to a lesser extent, exposure assessment, confounding and incomplete outcome data. Recruitment bias was low in three studies, probably low in 29 studies and high in another four, figure two. Risk of bias in exposure assessment was low in 17 studies, probably low in 8 and probably high in 11, figure two. Bias due to confounding was low risk in 30, probably low risk in one, probably high risk in a further four and high risk in one study. Incomplete outcome data was at low risk for bias in 18, probably low risk in ten, probably high risk in seven and high risk in one study. Risk for bias from selective reporting was low in 33 studies, probably low in two more and probably high in one. Supplemental table one explains reasons why a study was not rated at low risk of bias for the eight domains

considered. The quality and strength of evidence are described later for each separate category of exposure.

Overview of findings

Table 1 describes the exposure, the trimester where an association was sought and the direction of any association between exposures and fetal measurement. Regarding consistency of findings for second trimester measurements, an association was present between a maternal exposure in 16 of 88 (18%) comparisons made, and these associations were distributed evenly across all fetal measurements and all the exposures measured (table 1 and supplemental table 2). Third trimester measurements were associated with maternal exposures in 443 of the 96 comparisons made (46%) and were present for 15 of 26 (58%) analyses where BPD or HC were considered, 9 of 28 (32%) where FL was considered, 8 of 19 (42%) where AC was considered and 12 of 23 (52%) where EFW was considered (table 1 and supplemental table 2). Meta-analysis was not possible for maternal exposures to dietary factors, alcohol and other chemicals since either only one exposure was reported in the literature or if more than one study reported the same outcome, it was not valid or possible to pool the data. Meta-analysis was therefore only considered for ambient air exposures since comparable data seemed available (but not for other exposure categories), however the studies identified used a variety of methods to measure and report exposure (including units) meaning that meta analysis was not possible (table 2).

Maternal dietary exposures

Maternal plasma nutrient concentrations and fetal measurements

Vitamin D. FL was linked to maternal plasma 25(OH)D hydroxyl vitamin D in one study of 171 mothers aged <18 years (Young BE et al. 2012) and reduced exposure (i.e. ≤ 50 nmol/L) was associated with reduced FL (mean 0.15 z score) and humerus length (mean 0.18 z score) compared to higher exposure (i.e. >50 nmol/L). The associated reduction was limited to those whose mothers had both reduced plasma 25(OH)D and reduced dietary calcium

1 intake (i.e. <1050 mg/d)(Young BE et al. 2012); this study was at high risk for recruitment
2 bias and probably low risk for bias in two other domains of bias. Two studies from the
3 SWS(Mahon P et al. 2010; Ioannou C et al. 2012), one with probably low risk for bias in
4 confounding and probably high risk for bias due to incomplete outcome(Mahon P et al.
5 2010), used 3-D ultrasound technology to explore associations between maternal plasma
6 vitamin D status at 34 weeks gestation and fetal femur dimensions. There was no
7 association between maternal vitamin D and FL but there were associations with
8 characteristics of the femur. One study, which included data from 424 fetuses, found an
9 association between reduced maternal 25(OH)D (i.e. <25 nmol/L) and greater splaying of
10 the distal femur at 19 and 34 weeks(Mahon P et al. 2010). The second study(Ioannou C et
11 al. 2012) reported a weak positive association between maternal 25 (OH)D and proximal
12 metaphyseal diameter ($r=0.18$), but not estimated femur volume at 34 weeks gestation in
13 357 fetuses.

14 Vitamin E. Maternal plasma α -tocopherol (vitamin E) was positively linked to CRL at ten
15 weeks gestation in 766 mother-fetus pairs: the mean CRL was 42mm and 46 mm for the
16 lowest and highest quartiles, respectively(Turner SW et al. 2010). Maternal exposure was
17 not linked to fetal size later in gestation. The study was at probably low risk for bias in
18 incomplete outcome.

19 Fatty Acids. A Danish study related maternal whole blood fatty acid composition at 24
20 weeks gestation to fetal measurements made at 20 weeks gestation in 583 fetuses(Carlsen
21 K et al. 2013), and found a weak relationship between increasing n-3 PUFA and reducing FL
22 (but not HC or AC), the regression coefficient was -0.15 $p=0.02$.

23 Folate, vitamin B12 and homocysteine. A study from the Generation R cohort related
24 maternal and cord plasma concentrations of folate and vitamin B12, and also homocysteine
25 (associated with reduced birth weight and antioxidant properties) at 13 weeks gestation to
26 fetal weight gain and size at birth(Bergen et al. 2016). The main finding was that lower folate
27 and higher homocysteine in maternal and cord plasma were associated with a slowing of
28 fetal growth as pregnancy progressed (equivalent to 0.3 z score reduction in birth weight
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

between the upper and lower quartile groups). Unexpectedly, the highest quintile of maternal and cord plasma vitamin B12 was associated with lower birth weight compared to the lowest quintile (mean difference 43g for maternal plasma and 258g for cord plasma). A second Generation R study (Timmermans S et al. 2009) related timing of folate supplementation to EFW in 6365 fetuses and, consistent with the previously mentioned study (Bergen et al. 2016), found increased second trimester AC, third trimester AC and HC and EFW growth for those whose mothers started folic acid before or at the time of conception relative to no folic acid (mean EFW difference 0.10 z scores [0.02, 0.19]); individuals whose mothers who started folate supplementation in the first eight weeks had greater EFW growth in the second and third trimester (but no other fetal measurements) compared to no folic acid. A third Generation R study linked plasma folate concentration to fetal head growth and identified a small positive association equivalent to 3mm at birth per standard deviation increase in folate concentrations (Steenweg-de Graaff et al. 2017).

Maternal dietary intake and fetal measurements

Energy rich diet. A study from the Generation R cohort (Bouwland-Both MI et al. 2013) reported that an energy rich diet (i.e. rich in bread, nuts and margarine and therefore vitamins D and E), was associated with increased size in first trimester in 847 fetuses (mean increase in CRL 1.6mm relative to low energy rich diet); there was no association between diet and fetal measurements in later pregnancy. An observational study from Pakistan recruited 240 pregnant women (80 in each trimester, half of whom who fasted and half who did not fast during the month of Ramadan) and although mothers who fasted were 1kg lighter in the second and third trimester, there was no difference in growth in FL, BPD or EFW in any of the trimesters (Karateke et al. 2015); this study was at probably high risk for bias for confounding since socioeconomic status was not considered.

Mediterranean diet. A study from the Generation R cohort (Timmermans S et al. 2012) linked maternal Mediterranean diet to fetal measurements (n=3207), and individuals whose mothers were in the tertile with highest adherence to a "Mediterranean diet" (characterised

by higher intake of fruit, vegetables, fish, pasta and rice) had higher EFW at 20 and 30 weeks and increased AC at 30 weeks compared to those in the lowest adherence tertile (mean difference in z scores 0.11 for both measurements). The study was at probably high risk for bias in exposure assessment and probably low risk from incomplete data.

Milk. A further Generation R cohort paper described associations between maternal milk intake, as reported by food frequency questionnaire at 13 weeks gestation, and fetal measurements at 21 and 30 weeks gestation in 3405 mother-fetus pairs(Heppe DH et al. 2011b). There were associations between higher intakes and increased second trimester EFW and third trimester HC; with reference to ≤ 1 glass/day, 1-2 glasses/day was associated with 0.8mm increased HC and a 6g increased EFW. The study was at probably high risk for bias in exposure assessment and incomplete data.

Fish and seafood. A further Generation R study found no association between maternal fish and seafood consumption to fetal measurements at 21 and 30 weeks(Heppe DH et al. 2011a). A report from the EDEN mother-child cohort also found no association between maternal seafood intake and fetal measurements (Drouillet P et al. 2009). A second report from this cohort found no consistent evidence for mercury contamination of seafood to be associated with reduced second or third trimester fetal size in 691 mother-child pairs where maternal hair mercury was assessed (Drouillet-Pinard P et al. 2010). Both reports from the EDEN study were scored probably low risk for incomplete outcome since they reported significant associations between exposure and outcome for the subset whose mothers were overweight; this analysis was not pre-specified and the findings may be false positives.

Quality and strength of evidence

Collectively the quality of evidence linking maternal dietary exposures to fetal measurements was low since there was low risk of bias in population recruitment in all but one population, and often further risk from (questionnaire-based) exposure assessment and from incomplete outcomes. The strength of evidence was inadequate to relate exposure to fetal size and growth due to lack of replication for most exposures and differences in methodology between studies which did consider the same dietary exposure.

Ambient air pollution

Ten studies described associations between maternal air pollution exposure and fetal ultrasound scan measurement. Three studies were from the INMA cohort. Air pollutant exposures were: nitrogen dioxide (NO₂), particulate matter <10 microns (PM₁₀), benzene, sulphur dioxide (SO₂), carbon monoxide (CO) and ozone (O₃). Table 2 provides further details of the methodology used and magnitude of any association.

Nitrogen dioxide. Nine studies described the relationship between maternal NO₂ exposure and fetal measurement. Seven studies described associations between higher NO₂ exposures and reduced fetal size and/or growth(Aguilera I et al. 2010; Clemens et al. 2017; Iniguez C et al. 2012; Iniguez et al. 2016; Malmqvist et al. 2017; Ritz B et al. 2014; van den Hooven EH et al. 2012) and two studies reported no association(Carvalho M.A. et al. 2016; Hansen CA et al. 2008). The first of three studies from the INMA cohort, limited to 562 mothers in one of the recruitment centres (Sabadell), found increased NO₂ exposure was associated with a restriction in HC growth at weeks 12-20 and a reduction in BPD, intAC and EFW restriction at weeks 20-32 but only among mothers who spent <2 hours outdoors in non-residential areas per day(Aguilera I et al. 2010). The second report from the INMA cohort(Iniguez C et al. 2012) extended their previous results(Aguilera I et al. 2010) by relating cumulative NO₂ exposure during pregnancy to fetal size in a larger proportion of the cohort (785 mother). Here, the authors demonstrated an association for all mothers between increased maternal NO₂ exposure (i.e. 38 µg/m³) and a reduction in growth for BPD, AC and EFW between 20 and 34 weeks gestation; the authors concluded that NO₂ exposure before 20 weeks gestation was critical to the associations described. The third paper from the INMA cohort extended the previous findings(Iniguez C et al. 2012) to 2478 mothers(Iniguez et al. 2016) and the latter found evidence of faltering growth in the second (as well as the third) trimester associated with increasing NO₂ exposure. Two papers observed associations between increasing maternal NO₂ exposure and reduced fetal head

size, but no other fetal measurement(Clemens et al. 2017; Ritz B et al. 2014). One study reported increased maternal NO₂ exposure was associated with both reduced fetal head size and FL(van den Hooven EH et al. 2012) and another with both reduced FL and AC but not HC or BPD at 32-33 weeks gestation(Malmqvist et al. 2017). A small study of 366 mothers where NO₂ exposure was measured using personal passive samplers in each trimester found no association between exposure and third trimester EFW(Carvalho M.A. et al. 2016).

Fine particulates (PM₁₀). Three of the four papers which related maternal PM₁₀ exposure to fetal measurements found a link between increasing exposure and smaller measurements(Clemens et al. 2017; Hansen CA et al. 2008; van den Hooven EH et al. 2012). Where associations were present, they were found with reduced head size. The association with reduced head size was present only in the second trimester in one study(Hansen CA et al. 2008) and only after the second trimester in the other two studies(Clemens et al. 2017; van den Hooven EH et al. 2012). The study by Clemens *et al*(Clemens et al. 2017) found the association with PM₁₀ was also present for smaller particulates (PM_{2.5}) and also that the association was restricted to fetuses of non-smoking mothers. One study found an association between PM₁₀ exposure and reduced FL(Hansen CA et al. 2008) and a fourth found no association with any fetal measurement(Ritz B et al. 2014). The study by Hansen *et al* (Hansen CA et al. 2008) did not include socioeconomic status was rated probably high risk for bias in confounding.

Benzene. Two studies described an association between increased maternal benzene exposure and reduced head size(Aguilera I et al. 2010; Slama R et al. 2009). One study which was restricted to non-smoking mothers reported reduced second and third trimester fetal head size associated with increasing benzene exposure (equivalent to 2mm difference by 35 weeks gestation)(Slama R et al. 2009). The second study found no association between benzene and other fetal measurements including FL, AC and EFW, but found

reduced BPD growth between weeks 20-32 associated with benzene exposure among a subset of mothers who did not spend ≥ 15 hours at home per day(Aguilera I et al. 2010).

Other exposures. One study found an association between maternal ozone exposure in the first trimester and AC(Hansen CA et al. 2008) but two other studies found no association between ozone exposure and any fetal measurement(Carvalho M.A. et al. 2016; Ritz B et al. 2014). Only one study related SO_2 exposure to fetal measurements and reported associations between increased first trimester exposure and reduced BPD and AC(Hansen CA et al. 2008). One study explored the relationship between CO and fetal measurements and found no associations(Ritz B et al. 2014).

Quality and strength of evidence

Collectively the ten well-designed well-conducted studies provided a high quality of evidence linking maternal exposure to NO_2 and $\text{PM}_{2.5}$ to reduced third trimester fetal measurements (especially head size). The “default” for human studies having moderate quality of evidence was upgraded on account of evidence of dose response between NO_2 and $\text{PM}_{2.5}$ exposures and reduced fetal head size. Many studies were at probably low risk for bias in recruitment and exposure assessment. There was sufficient evidence to link NO_2 and $\text{PM}_{2.5}$ to reduced fetal head size since findings were consistent across a number of studies, reporting different concentrations of exposures, using different design and in different countries. The evidence linking other air pollution exposures to fetal measurements was of high quality but there was inadequate evidence of an association due to inconsistent findings between studies (e.g. benzene) and not all exposures were measured in all studies, e.g. ozone, carbon monoxide.

Maternal alcohol intake

One paper from the Generation R cohort assessed maternal alcohol intake in the first, second and third trimesters for 7333 mother-fetal pairs(Bakker R et al. 2010). In the longitudinal analysis, the fetuses of mothers who continued to consume alcohol (typically <1

1 drink a week) had a small increase in EFW growth (0.6g per week). In cross-sectional
2 analyses, there was no association between second and third trimester intake and fetal HC,
3 AC, FL or EFW. There was no evidence of a dose-response effect of alcohol on fetal
4 measurements. A second publication which excluded mothers who smoked obtained
5 routinely collected ultrasound measurements between 18 and 41 weeks gestation and
6 reported reduced HC:AC during the pregnancy for fetuses whose mothers who continued to
7 drink compared to those who quit (magnitude not described)(Handmaker NS et al. 2006).
8 There was reduced cerebellar growth, but no difference in HC, AC and FL, among fetuses
9 exposed to persistent maternal alcohol intake compared to those whose mothers who quit
10 drinking during pregnancy. In this cohort(Handmaker NS et al. 2006), 40% of mothers were
11 marijuana users and 20% used amphetamines with the latter being associated with
12 increased HC:AC growth. A third study (a pilot study) screened 6745 Ukrainian mothers in
13 early pregnancy and identified 84 moderate-to-heavy drinkers and 82 abstinent mothers(Kfir
14 M et al. 2009); moderate-to-heavy drinking was a reported average of 30mls ethanol (or
15 three UK “units”) daily at conception and 4mls daily after knowingly being pregnant (including
16 binges with an average of 66mls ethanol/day). FL was reduced in the second trimester
17 among alcohol exposed fetuses (mean 53rd centile versus 65th centile for controls) but not in
18 the third trimester. Third trimester BPD was reduced in exposed fetuses when compared to
19 controls (54th centile versus 70th centile, respectively).
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 Collectively these studies were of low quality due to bias in recruitment, use of reported
45 alcohol intake and incomplete follow up and within this limited literature there was
46 inadequate evidence identified to link maternal alcohol intake to fetal measurements.
47
48
49
50
51
52

53 **Other exposures**

54 *Occupational exposures.* A study of occupational exposures (by questionnaire) in 4680
55 pregnant mothers related FL, HC and EFW to exposures including phthalates, pesticides,
56 polyaromatic hydrocarbons and alkylphenolic compounds(Snijder CA et al. 2012b). There
57
58
59
60
61
62
63
64
65

1 were associations of small magnitude seen between some exposures and some fetal
2 measurements, most consistently for phthalates whose exposure was associated with
3 reduced EFW and fetal length (approximately 1% of a z score per week).
4

5
6 *Bisphenol A.* A second Generation R study measured maternal urinary Bisphenol A (BPA) in
7 the first, second and third trimesters and related these to fetal growth in 419 pregnancies
8 (Snijder CA et al. 2013). In 80 mothers (i.e. 19% of the study population) urine samples
9 were obtained in each trimester there was reduced growth EFW and HC per unit increase in
10 BPA (typically 2% reduction per week per unit increase in BPA exposure), but this finding
11 was not replicated in models which included mothers where exposure data were not
12 complete nor in a report from the INMA cohorts where there was no association between
13 maternal first and third trimester BPA and fetal measurements (488 mothers)(Casas et al.
14 2016).
15

16
17 *Phenols.* A study from the EDEN cohort related maternal exposure to nine phenol-based
18 chemicals to fetal size and growth throughout pregnancy and found evidence that one
19 (triclosan) was linked to minor reductions in fetal size in the third trimester but at no other
20 time(Philippat et al. 2014).
21

22
23 *Phthalates.* Two studies related maternal phthalate exposure to fetal size and growth and
24 report apparently contrasting findings. The first study, again restricted to male offspring in
25 the EDEN cohort, measured 11 phthalate molecules in maternal urine and found a negative
26 relationship between phthalate metabolites and EFW (10-15g per quartile increase in
27 phthalate exposure) but also a positive association between one molecule
28 (monocarboxyisononyl) and FL in the second and third trimesters(Botton et al. 2016) (which
29 is consistent with the previously mentioned study(Casas et al. 2016)). A study from the INMA
30 cohorts found association of small magnitude between increased concentrations of two of
31 eight phthalate molecules in maternal urine and reduced growth HC in the first, and FL
32 second half of pregnancy(Casas et al. 2016). Both cohorts only measured one common
33 phthalate molecule (monoethyl phthalate) so the results are not necessarily inconsistent.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Polybrominated diphenyl ethers (PBDEs). A further evaluation of the INMA cohorts measured PBDEs in maternal and cord concentrations and related these to fetal measurements in 670 mothers (Lopez-Espinosa et al. 2015); a doubling of PBDE exposure was associated with <5% reduced growth in AC, EFW and BPD (but not FL) between 20-34 weeks (but not 12-20 weeks), although associations were not always present for both maternal and cord blood concentrations.

Organochlorine compounds. The final study identified from 2369 the INMA cohorts related maternal organochlorine compounds in maternal plasma at 12 weeks gestation and cord blood to fetal growth in early, mid and late pregnancy (Lopez-Espinosa et al. 2016); a doubling of exposures to polychlorinated biphenyls (PCB) -138, -153 and -180 were associated with a 2-4% reduction in femur length growth and a doubling of PCB-138 was associated with a 2% reduction in EFW growth between gestational weeks 20-34. There were inverse associations between cord Hexachlorobenzene and reduced AC growth between 0 and 20 weeks. Five of 120 associations in this analysis achieved significance.

Lithium in drinking water. The northern Argentinian Andes has variable concentrations of lithium in its drinking water and a study of 194 mothers observed no significant relationships between maternal plasma and urinary lithium and second and third trimester BPD, HC, AC, FL and EFW (Harari et al. 2015).

Collectively the quality of evidence for these studies was low since although objective measurements of exposure were made in all but one study, multiple comparisons were made within studies increasing the chance of false positive findings, findings (when replicated) were not consistent and risk of bias from incomplete follow up was probably high or high in four studies. Additionally, data were limited only to male offspring in two studies (Botton et al. 2016; Philippat et al. 2014). In conclusion, there was inadequate strength of evidence linking chemicals to fetal size since the quality of evidence was low, many different chemicals were studied and where associations with fetal size were present, these were not replicated elsewhere.

DISCUSSION

This systematic review was designed to describe the literature associating maternal exposures to fetal size being small for gestational age. The papers identified were all published since 2006, indicating that this is a relatively new literature. The first major finding was that maternal exposures to increased ambient air NO₂ and PM_{2.5} were consistently associated with reduced third trimester fetal head size and we judge that the literature presents sufficient evidence of toxicity in this context. The second major finding was that the strength of evidence was inadequate for all the other exposures considered. The third notable finding was that, where associations were present, they were more commonly seen in the third compared to the second trimester and that reduced third trimester head size was more commonly associated with potentially adverse maternal exposures compared to other fetal measurements. Together these findings suggest that public health measures are urgently required to minimise pregnant mother's exposure to NO₂ and PM_{2.5}, and more high-quality research is required to better understand the relationship between other (modifiable) maternal exposures and fetal measurements.

The fetus has traditionally been thought to have a privileged position, where it was protected from the adverse effects of environmental exposures by the maternal-placental "unit", but associations between reduced birth weight and many maternal exposures argue against this paradigm. The fetal origins hypothesis (Barker 1995a) speculated that maternal exposures in mid pregnancy were relevant to birth weight and risk for subsequent increased risk for non-communicable diseases. Since we find evidence of reduced second trimester fetal size and some maternal exposures, and our findings suggest that maternal exposures in early pregnancy are also relevant.

There were some associations between exposure and fetal size which were mostly consistent across different populations, whilst other associations were less consistently seen, or were even counterintuitive. An example of consistency was the inverse association

1 between NO₂ exposure and third trimester fetal size, which was seen in all seven studies
2 which explored this link (two studies measured NO₂ but not third trimester head size
3
4 Carvalho M.A. et al. 2016; Hansen CA et al. 2008). In contrast, there were some instances
5
6 where an apparently harmful exposure was associated with increased fetal size (e.g. current
7
8 PM₁₀ exposure and first trimester length and second trimester weight(van den Hooven EH et
9
10 al. 2012), one phthalate molecule and increased third trimester fetal length(Botton et al.
11
12 2016), increased maternal and cord plasma vitamin B12 concentrations and reduced birth
13
14 weight(Bergen et al. 2016), maternal alcohol intake and increased third trimester
15
16 EFW(Bakker R et al. 2010)); although these associations may challenge the paradigm that
17
18 “harmful” exposures invariably cause reduced fetal growth, these may be false positive
19
20 findings.
21
22
23
24

25 A further example of inconsistency comes from the ambient air exposure literature where
26
27 two studies found that the magnitude of association between NO₂ and fetal size was only
28
29 present(Ritz B et al. 2014), or was greater(Iniguez et al. 2016), among mothers who smoked
30
31 whereas a second study only found an association between NO₂ and fetal head size among
32
33 non-smokers(Clemens et al. 2017). Our review also identified many associations of
34
35 borderline significance and instances of multiple testing, and these increase the risk of false
36
37 positive results. Given that the body of literature reviewed is relatively young, it is not
38
39 unexpected that there is a spectrum of consistency/inconsistency between maternal
40
41 exposure and fetal size and further research activity is required to replicate some of the
42
43 apparently inconsistent associations.
44
45
46
47

48 There are consistencies and inconsistencies in the literature describing associations
49
50 between maternal exposures and birth weight and these exposures and fetal size. Maternal
51
52 exposures to poor quality ambient air is associated with reduced birth weight (Maisonet et al.
53
54 2004; Shah PS. Balkhair T. Knowledge Synthesis Group on Determinants of Preterm/LBW
55
56 births 2011) and we confirm this association is already present in fetal life. In contrast, many
57
58 maternal dietary exposures are associated with reduced birth weight (Gresham E et al.
59
60
61
62
63
64
65

2014) but there were no consistent associations between maternal dietary exposures and fetal size in the literature we reviewed. Similarly, maternal exposure to chemicals such as bisphenol A (Perera F and Herbstman J 2011) is associated with reduced birth weight but bisphenol A was associated with reduced fetal size in one study (Casas et al. 2016) but not a second (Snijder CA et al. 2013). The inconsistent findings between size before and after birth in the context of maternal exposures may reflect a smaller literature describing antenatal size, the challenges of accurately measuring fetal size or theoretically may be due to associations only becoming detectable towards the very end of pregnancy.

Maternal alcohol ingestion during pregnancy is an established risk factor for reduced birth weight (Henderson J et al. 2007), but the literature was not adequate to determine when the relationship between alcohol ingestion and fetal growth failure begins. The studies identified were at high risk of bias and collectively provided inadequate evidence of toxicity. This absence of evidence does not mean that any change is required to current guidelines which recommend that pregnant mothers should abstain from drinking alcohol.

A weakness of the present literature is that all the evidence comes from observational studies where, in addition to the risk of false positive findings previously discussed, some of the results may be influenced by confounding factors and also the results are based on populations who are not necessarily representative of the general population. Drop out from cohort studies also may contribute bias, for example a study which related fetal size to risk for later asthma found a relationship of greater magnitude when using questionnaire reported asthma outcome (available in 39% of the cohort) compared to using routinely acquired asthma outcome (available in 88%) (Turner et al. 2018).

One further limitation of the literature is that whilst there are some maternal exposures linked to fetal size in three or more studies, for example smoking (Abraham et al. 2017) and air pollution (the present review), many exposures are linked to fetal size in only one or two studies (for example chemicals and dietary nutrients) and this leaves insufficient evidence

1 upon which to form an opinion. A second potential limitation to the literature is that fetal
2 measurements and exposures were made at different gestational age, although this
3
4 limitation applies to the different studies in the literature linking post natal exposures and
5
6 post natal outcomes. Fetal size is driven by gestational age, and fetal size was determined
7
8 over a range of gestations within and between different studies. Gestational age was
9
10 included as a covariate in all but one study (Karateke et al. 2015) and this minimises the
11
12 potential for differences in gestational age at assessment confounding associations with
13
14 maternal exposure. Differences between studies in the gestational age when fetal size and
15
16 maternal exposure are likely to weaken and not strengthen the associations (where present)
17
18 between maternal exposure and fetal size.
19
20
21

22
23 A further limitation to the literature was that different methods were used to measure
24
25 exposures, even where the same exposure (NO₂ for example) was measured in several
26
27 studies. This meant that meta-analysis was not possible for single exposures. Another
28
29 limitation to the literature is that there are some areas where the evidence is of a poor
30
31 quality, e.g. maternal alcohol exposure, which is an increasing public health concern due to
32
33 increasing awareness of fetal alcohol syndrome(Popova et al. 2017). Finally, there are
34
35 several maternal exposures including recreational drug use (Behnke M et al. 2013),
36
37 pesticides(Sathyanarayana et al. 2010), proximity to land fill sites(Elliott et al. 2001) and
38
39 electromagnetic fields (de Vocht F and Lee B 2014) which are linked to reduced birth weight
40
41 but where there are no studies associating such exposures to fetal measurements.
42
43
44
45

46 Pregnant mothers are constantly exposed to multiple environmental exposures and
47
48 identifying which single exposure may be important to fetal size is a challenge, for example
49
50 increased maternal alcohol intake may be associated with smoking, use of illicit drugs and
51
52 poor diet. Furthermore, any reduction in fetal size associated with tobacco and alcohol
53
54 consumption may be modified by increased size associated with amphetamine intake
55
56 (Handmaker NS et al. 2006). Only intervention studies are able to infer causality, but these
57
58 are practically and ethically challenging to do in the context of potentially harmful maternal
59
60
61
62
63
64
65

exposures during pregnancy and the evidence base may remain dependent on observational studies.

The mechanism(s) where environmental exposures may cause fetal growth failure (or acceleration) are not well understood but likely to be multiple. The mechanism for maternal diet influencing fetal size assumes that macro and micro nutrients cross the placenta, and maternal dietary fat intake during the third trimester has been associated with cord blood lipid concentrations (Elias SL and Innis SM 2001) but the fetus is not necessarily exposed to all maternal plasma nutrients, for example triglycerides do not cross the placenta(Herrera 2002). The mechanism for ambient air exposures leading to fetal growth failure is not understood but ultrafine particles (which correlate with PM₁₀ and NO₂ concentrations) are able to directly enter the maternal circulation via the alveoli (Geiser M et al. 2005) and placenta (Wick P et al. 2010), and thus theoretically enter the fetal circulation where they may be pathogenic by causing vascular inflammation(Gojova A et al. 2007).

In summary, there is an emerging literature which suggests that some maternal exposures during early pregnancy are associated with changes in fetal size which are apparent by the second and third trimester. Evidence from randomised controlled trials where maternal exposures are modified are unlikely due to practical and ethical issues, and the evidence base is likely to arise from observational studies. The literature is mostly based on Western populations and is dominated by three cohort studies and replication in other populations of the associations described is required.

ACKNOWLEDGEMENTS

The authors thank Dr Juleen Lam for her advice on the use of the Navigation guide and for her helpful comments on a draft of the submitted manuscript.

REFERENCES

- 1
2
3 Abraham, M.; Alramadhan, S.; Iniguez, C.; Duijts, L.; Jaddoe, V.W.V.; Den Dekker, H.T. et
4 al. A systematic review of maternal smoking during pregnancy and fetal measurements
5 with meta-analysis. PLoS ONE 12:e0170946; 2017.
6
- 7 Aguilera I; Garcia-Esteban R; Iniguez C; Nieuwenhuijsen MJ; Rodriguez A; Paez M et al.
8 Prenatal exposure to traffic-related air pollution and ultrasound measures of fetal growth
9 in the INMA Sabadell cohort. Environ Health Perspect 118:705-11; 2010.
10
- 11 Alkandari, F.; Ellahi, A.; Aucott, L.; Devereux, G.; Turner, S. Fetal ultrasound measurements
12 and associations with postnatal outcomes in infancy and childhood: a systematic review
13 of an emerging literature. J Epidemiol Comm Health 69:41-48; 2015.
14
- 15 Bakker R; Pluimgraaff LE; Steegers EA; Raat H; Tiemeier H; Hofman A et al. Associations of
16 light and moderate maternal alcohol consumption with fetal growth characteristics in
17 different periods of pregnancy: the Generation R Study. Int J Epidemiol 39:777-89;
18 2010.
19
- 20 Barker, D.J. Fetal origins of coronary heart disease. BMJ 311:171-4; 1995.
21
- 22 Bateson, P.; Barker, D.; Clutton-Brock, T.; Deb, D.; D'Udine, B.; Foley, R.A. et al.
23 Developmental plasticity and human health. Nature 430:419-21; 2004.
24
- 25 Behnke M; Smith VC; Committee on Substance Abuse; Committee on Fetus and Newborn.
26 Prenatal substance abuse: short- and long-term effects on the exposed fetus. Pediatrics
27 131:e1009-24; 2013.
28
- 29 Bergen, N.E.; Schalekamp-Timmermans, S.; Jaddoe, V.W.V.; Hofman, A.; Lindemans, J.;
30 Russcher, H. et al. Maternal and Neonatal Markers of the Homocysteine Pathway and
31 Fetal Growth: The Generation R Study. Paediatr Perinat Epidemiol 30:386-96; 2016.
32
- 33 Botton, J.; Philippat, C.; Calafat, A.M.; Carles, S.; Charles, M.; Slama, R. et al. Phthalate
34 pregnancy exposure and male offspring growth from the intra-uterine period to five
35 years of age. Environ Res 151:601-9; 2016.
36
- 37 Bouwland-Both MI; Steegers-Theunissen RP; Vujkovic M; Lesaffre EM; Mook-Kanamori DO;
38 Hofman A et al. A periconceptional energy-rich dietary pattern is associated with early
39 fetal growth: the Generation R study. BJOG: An International Journal of Obstetrics &
40 Gynaecology 120:435-45; 2013.
41
- 42 Carlsen K; Pedersen L; Bonnelykke K; Stark KD; Lauritzen L; Bisgaard H. Association
43 between whole-blood polyunsaturated fatty acids in pregnant women and early fetal
44 weight. Eur J Clin Nutr 67:978-83; 2013.
45
- 46 Carvalho M.A.; Bernardes L.S.; Hettfleisch K.; Pastro L.D.M.; Vieira S.E.; Saldiva S.R.D.M.
47 et al. Associations of maternal personal exposure to air pollution on fetal weight and
48 fetoplacental Doppler: A prospective cohort study. Reproductive Toxicology 62:9-17;
49 2016.
50
- 51 Casas, M.; Valvi, D.; Ballesteros-Gomez, A.; Gascon, M.; Fernandez, M.F.; Garcia-Esteban,
52 R. et al. Exposure to Bisphenol A and Phthalates during Pregnancy and Ultrasound
53
54
55
56
57
58
59
60
61
62
63
64
65

- Measures of Fetal Growth in the INMA-Sabadell Cohort. *Environ Health Perspect* 124:521-8; 2016.
- Clemens, T.; Turner, S.; Dibben, C. Maternal exposure to ambient air pollution and fetal growth in North-East Scotland: A population-based study using routine ultrasound scans *Environ Int* 107:216-226; 2017.
- de Vocht F and Lee B. Residential proximity to electromagnetic field sources and birth weight: Minimizing residual confounding using multiple imputation and propensity score matching. *Environ Int* 69:51-7; 2014.
- Drouillet P; Kaminski M; De Lauzon-Guillain B; Forhan A; Ducimetiere P; Schweitzer M et al. Association between maternal seafood consumption before pregnancy and fetal growth: evidence for an association in overweight women. The EDEN mother-child cohort. *Paediatr Perinat Epidemiol* 23:76-86; 2009.
- Drouillet-Pinard P; Huel G; Slama R; Forhan A; Sahuquillo J; Goua V et al. Prenatal mercury contamination: relationship with maternal seafood consumption during pregnancy and fetal growth in the 'EDEN mother-child' cohort. *Br J Nutr* 104:1096-100; 2010.
- Elias SL and Innis SM. Infant plasma trans, n-6, and n-3 fatty acids and conjugated linoleic acids are related to maternal plasma fatty acids, length of gestation, and birth weight and length. *Am J Clin Nutr* 73:807-14; 2001.
- Elliott, P.; Briggs, D.; Morris, S.; de Hoogh, C.; Hurt, C.; Jensen, T.K. et al. Risk of adverse birth outcomes in populations living near landfill sites. *BMJ* 323:363-8; 2001.
- Geiser M; Rothen-Rutishauser B; Kapp N; Schurch S; Kreyling W; Schulz H et al. Ultrafine particles cross cellular membranes by nonphagocytic mechanisms in lungs and in cultured cells. *Environ Health Perspect* 113:1555-60; 2005.
- Gluckman, P.D.; Hanson, M.A.; Low, F.M. The role of developmental plasticity and epigenetics in human health. *Birth Defects Research. Part C, Embryo Today: Reviews* 93:12-8; 2011.
- Gluckman, P.D.; Hanson, M.; Spencer, H.G. Predictive adaptive responses and human evolution. *Trends Ecol Evol* 20:527-33; 2005.
- Gojova A; Guo B; Kota RS; Rutledge JC; Kennedy IM; Barakat AI. Induction of inflammation in vascular endothelial cells by metal oxide nanoparticles: effect of particle composition. *Environ Health Perspect* 115:403-9; 2007.
- Gresham E; Byles JE; Bisquera A; Hure AJ. Effects of dietary interventions on neonatal and infant outcomes: a systematic review and meta-analysis. *Am J Clin Nutr* 100:1298-321; 2014.
- Guxens M; Ballester F; Espada M; Fernandez MF; Grimalt JO; Ibarluzea J et al. Cohort Profile: the INMA--Infancia y Medio Ambiente--(Environment and Childhood) Project. *Int J Epidemiol* 41:930-40; 2012.
- Hadlock, F.P.; Harrist, R.B.; Carpenter, R.J.; Deter, R.L.; Park, S.K. Sonographic estimation of fetal weight. The value of femur length in addition to head and abdomen measurements. *Radiology* 150:535-40; 1984.

- Hales CN and Barker DJ. Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis. *Diabetologia* 35:595-601; 1992.
- Handmaker NS; Rayburn WF; Meng C; Bell JB; Rayburn BB; Rappaport VJ. Impact of alcohol exposure after pregnancy recognition on ultrasonographic fetal growth measures. *Alcoholism: Clinical & Experimental Research* 30:892-8; 2006.
- Hansen CA; Barnett AG; Pritchard G. The effect of ambient air pollution during early pregnancy on fetal ultrasonic measurements during mid-pregnancy. *Environ Health Perspect* 116:362-9; 2008.
- Harari, F.; Langeen, M.; Casimiro, E.; Bottai, M.; Palm, B.; Nordqvist, H. et al. Environmental exposure to lithium during pregnancy and fetal size: a longitudinal study in the Argentinean Andes. *Environ Int* 77:48-54; 2015.
- Henderson J; Kesmodel U; Gray R. Systematic review of the fetal effects of prenatal binge-drinking. *Journal of Epidemiology & Community Health* 61:1069-73; 2007.
- Heppe DH; Steegers EA; Timmermans S; Breeijen Hd; Tiemeier H; Hofman A et al. Maternal fish consumption, fetal growth and the risks of neonatal complications: the Generation R Study. *Br J Nutr* 105:938-49; 2011a.
- Heppe DH; van Dam RM; Willemsen SP; den Breeijen H; Raat H; Hofman A et al. Maternal milk consumption, fetal growth, and the risks of neonatal complications: the Generation R Study. *Am J Clin Nutr* 94:501-9; 2011b.
- Herrera, E. Implications of dietary fatty acids during pregnancy on placental, fetal and postnatal development--a review. *Placenta* 23:S9-19; 2002.
- Iniguez C; Ballester F; Estarlich M; Esplugues A; Murcia M; Llop S et al. Prenatal exposure to traffic-related air pollution and fetal growth in a cohort of pregnant women. *Occupational & Environmental Medicine* 69:736-44; 2012.
- Iniguez, C.; Esplugues, A.; Sunyer, J.; Basterrechea, M.; Fernandez-Somoano, A.; Costa, O. et al. Prenatal Exposure to NO₂ and Ultrasound Measures of Fetal Growth in the Spanish INMA Cohort. *Environ Health Perspect* 124:235-42; 2016.
- Ioannou C; Javaid MK; Mahon P; Yaqub MK; Harvey NC; Godfrey KM et al. The effect of maternal vitamin D concentration on fetal bone. *Journal of Clinical Endocrinology & Metabolism* 97:E2070-7; 2012.
- Johnson, P.I.; Sutton, P.; Atchley, D.S.; Koustas, E.; Lam, J.; Sen, S. et al. The Navigation Guide - evidence-based medicine meets environmental health: systematic review of human evidence for PFOA effects on fetal growth. *Environ Health Perspect* 122:1028-39; 2014.
- Karateke, A.; Kaplanoglu, M.; Avci, F.; Kurt, R.K.; Baloglu, A. The effect of Ramadan fasting on fetal development.. *Pakistan Journal of Medical Sciences* 31:1295-9; 2015.
- Kfir M; Yevtushok L; Onishchenko S; Wertelecki W; Bakhireva L; Chambers CD et al. Can prenatal ultrasound detect the effects of in-utero alcohol exposure? A pilot study. *Ultrasound in Obstetrics & Gynecology* 33:683-9; 2009.

- 1 Lopez-Espinosa, M.; Costa, O.; Vizcaino, E.; Murcia, M.; Fernandez-Somoano, A.; Iniguez,
2 C. et al. Prenatal Exposure to Polybrominated Flame Retardants and Fetal Growth in
3 the INMA Cohort (Spain). *Environ Sci Technol* 49:10108-16; 2015.
- 4 Lopez-Espinosa, M.; Murcia, M.; Iniguez, C.; Vizcaino, E.; Costa, O.; Fernandez-Somoano,
5 A. et al. Organochlorine Compounds and Ultrasound Measurements of Fetal Growth in
6 the INMA Cohort (Spain). *Environ Health Perspect* 124:157-163; 2016
- 7
8 Mahon P; Harvey N; Crozier S; Inskip H; Robinson S; Arden N et al. Low maternal vitamin D
9 status and fetal bone development: cohort study. *Journal of Bone & Mineral Research*
10 25:14-9; 2010.
- 11
12
13 Maisonet, M.; Correa, A.; Misra, D.; Jaakkola, J.J. A review of the literature on the effects of
14 ambient air pollution on fetal growth. *Environ Res* 95:106-15; 2004.
- 15
16
17 Malmqvist, E.; Liew, Z.; Kallen, K.; Rignell-Hydbom, A.; Rittner, R.; Rylander, L. et al. Fetal
18 growth and air pollution - A study on ultrasound and birth measures. *Environ Res*
19 152:73-80; 2017.
- 20
21
22 Pereira, P.P.d.S.; Da Mata, F.A.F.; Figueiredo, A.C.G.; de Andrade, K.R.C.; Pereira, M.G.
23 Maternal Active Smoking During Pregnancy and Low Birth Weight in the Americas: A
24 Systematic Review and Meta-analysis. *Nicotine Tobacco Res* 19:497-505; 2017.
- 25
26 Perera F and Herbstman J. Prenatal environmental exposures, epigenetics, and disease.
27 *Reproductive Toxicology* 31:363-73; 2011.
- 28
29
30 Philippat, C.; Botton, J.; Calafat, A.M.; Ye, X.; Charles, M.; Slama, R. et al. Prenatal
31 exposure to phenols and growth in boys.. *Epidemiology* 25:625-35; 2014.
- 32
33
34 Popova, S.; Lange, S.; Probst, C.; Gmel, G.; Rehm, J. Estimation of national, regional, and
35 global prevalence of alcohol use during pregnancy and fetal alcohol syndrome: a
36 systematic review and meta-analysis.. *The Lancet Global Health* 5:e290-9; 2017.
- 37
38
39 Ritz B; Qiu J; Lee PC; Lurmann F; Penfold B; Erin Weiss R et al. Prenatal air pollution
40 exposure and ultrasound measures of fetal growth in Los Angeles, California. *Environ*
41 *Res* 130:7-13; 2014.
- 42
43
44 Sathyanarayana, S.; Basso, O.; Karr, C.J.; Lozano, P.; Alavanja, M.; Sandler, D.P. et al.
45 Maternal pesticide use and birth weight in the agricultural health study. *J Agromed*
46 15:127-36; 2010.
- 47
48
49 Shah PS. Balkhair T. Knowledge Synthesis Group on Determinants of Preterm/LBW births.
50 Air pollution and birth outcomes: a systematic review. *Environ Int* 37:498-516; 2011.
- 51
52
53 Shaheen SO; Sterne JA; Montgomery SM; Azima H. Birth weight, body mass index and
54 asthma in young adults. *Thorax* 54:396-402; 1999.
- 55
56
57 Slama R; Thiebaugeorges O; Goua V; Aussel L; Sacco P; Bohet A et al. Maternal personal
58 exposure to airborne benzene and intrauterine growth. *Environ Health Perspect*
59 117:1313-21; 2009.
- 60
61
62
63
64
65

- 1 Snijder CA; Brand T; Jaddoe V; Hofman A; Mackenbach JP; Steegers EA et al. Physically
2 demanding work, fetal growth and the risk of adverse birth outcomes. The Generation R
3 Study. Occupational & Environmental Medicine 69:543-50; 2012a.
- 4 Snijder CA; Heederik D; Pierik FH; Hofman A; Jaddoe VW; Koch HM et al. Fetal growth and
5 prenatal exposure to bisphenol A: the generation R study. Environ Health Perspect
6 121:393-8; 2013.
- 7
8
9 Snijder CA; Roeleveld N; Te Velde E; Steegers EA; Raat H; Hofman A et al. Occupational
10 exposure to chemicals and fetal growth: the Generation R Study. Human Reproduction
11 27:910-20; 2012b.
- 12
13 Steenweg-de Graaff, J.; Roza, S.J.; Walstra, A.N.; El Marroun, H.; Steegers, E.A.P.; Jaddoe,
14 V.W.V. et al. Associations of maternal folic acid supplementation and folate
15 concentrations during pregnancy with foetal and child head growth: the Generation R
16 Study.. Eur J Nutr 56:65-75; 2017.
- 17
18
19 Timmermans S; Jaddoe VW; Hofman A; Steegers-Theunissen RP; Steegers EA.
20 Periconception folic acid supplementation, fetal growth and the risks of low birth weight
21 and preterm birth: the Generation R Study. Br J Nutr 102:777-85; 2009.
- 22
23
24 Timmermans S; Steegers-Theunissen RP; Vujkovic M; den Breeijen H; Russcher H;
25 Lindemans J et al. The Mediterranean diet and fetal size parameters: the Generation R
26 Study. Br J Nutr 108:1399-409; 2012.
- 27
28
29 Turner SW; Campbell D; Smith N; Craig LC; McNeill G; Forbes SH et al. Associations
30 between fetal size, maternal {alpha}-tocopherol and childhood asthma. Thorax 65:391-
31 7; 2010.
- 32
33
34 Turner, S.; Fielding, S.; Devereux, G. First trimester fetal size and prescribed asthma
35 medication at 15 years of age. European Respiratory Journal 51; 2018.
- 36
37 van den Hooven EH; Pierik FH; de Kluizenaar Y; Willemsen SP; Hofman A; van Ratingen
38 SW et al. Air pollution exposure during pregnancy, ultrasound measures of fetal growth,
39 and adverse birth outcomes: a prospective cohort study. Environ Health Perspect
40 120:150-6; 2012.
- 41
42
43 Wick P; Malek A; Manser P; Meili D; Maeder-Althaus X; Diener L et al. Barrier capacity of
44 human placenta for nanosized materials. Environ Health Perspect 118:432-6; 2010.
- 45
46
47 Young BE; McNanley TJ; Cooper EM; McIntyre AW; Witter F; Harris ZL et al. Maternal
48 vitamin D status and calcium intake interact to affect fetal skeletal growth in utero in
49 pregnant adolescents. Am J Clin Nutr 95:1103-12; 2012.
- 50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

FIGURE LEGENDS

Figure 1. CONSORT style diagram showing how the papers included in the review were identified.

Figure two. Bar chart summarising the risk of bias for each of the eight domains considered across the 36 papers included in this systematic review.

Figure three. This figure shows the risk of bias in each of eight domains considered for each of the 36 studies included in this systematic review.

Table 1. Summary of results from the papers identified in the systematic review. HC=head circumference, BPD=biparietal diameter, AC=abdominal circumference, MAD=mean abdominal circumference, EFW=estimated fetal weight. X=no association present, ↑exposure associated with increased fetal measurement, ↓ exposure associated with reduced fetal measurement. NO₂= nitrogen dioxide, CO=carbon monoxide, PM₁₀=particulates with diameter less than 10 microns, BTEX= aromatic hydrocarbons benzene, toluene, ethylbenzene, m/p-xylene, and o-xylene, SO₂=sulphur dioxide, O₃=ozone, PUFA=poly unsaturated fatty acids, PAH=polycyclic aromatic hydrocarbons.

	Exposure	Trimester when exposure was measured*	Second trimester				Third trimester			
			HC/BPD	FL	AC/MAD	EFW	HC/BPD	FL	AC/MAD	EFW
Drouillet (Drouillet P et al. 2009) (seafood)	Diet	P and 3	x	x	x		x	x	x	
Timmermans (Timmermans S et al. 2009)(Folic acid supplementation)		P and 1	x	x	↑	x	↑↑	x	↑↑	↑
Mahon(Mahon P et al. 2010) (Vit D)		3 (week 34)		x				x		
Turner(Turner SW et al. 2010) (Vit E)		1 (week 12)	x	x						
Drouillet-Pinard(Drouillet-Pinard P et al. 2010)(mercury)		P and 3	x	x	x		x	x	x	
Heppe(Heppe DH et al. 2011b)(milk)		1 (week 15)	x	x		↑	↑	x		x
Heppe(Heppe DH et al. 2011a) (fish)		1 (week 15)	x	x	x	x	x	x	x	x
Ioannou(Ioannou C et al. 2012)(Vit D)		3 (week 34)						↑§		
Young(Young BE et al. 2012) (Vit D and calcium)		3						↑		
Timmermans(Timmermans S et al. 2012)(Mediterranean diet)		1 (week 15)	x	x	x	↑	x	x	↑	↑
Carlsen(Carlsen K et al. 2013)(PUFA)		2 (week 24)	x	↓	x	x				
Bouwland-Both(Bouwland-Both MI et al. 2013) (energy rich diet)		1 (week 15)								x
Karateke(Karateke et al. 2015) (fasting)		1, 2 and 3	x‡	x‡		x‡	x‡	x‡		x‡
Bergen(Bergen et al. 2016) (Homocysteine)		1 (week 13)	x	x		x	↓	↓		↓

Steenweg(Steenweg-de Graaff et al. 2017) (folic acid)		1 (week 13)						↑ δ			
Hansen (Hansen CA et al. 2008)	Ambient air pollution¥	1 (first 120 days)	↓	↓	↓						
Slama (Slama R et al. 2009)		2 (week 27)	↓					↓			
Aguilera(Aguilera I et al. 2010)		1 (weeks 1-12) 2 (weeks 12-20) 3 (weeks 20-32)	↓ ψ	x	x	x		↓ ψ	x	↓ ψ	↓ ψ
Iniguez (Iniguez C et al. 2012)		1 (weeks 1-12) 2 (weeks 12-20) 3 (weeks 20-32, 32-term)	x	x	x	x		↓ ψ	x	↓ ψ	↓ ψ
van den Hooven(van den Hooven EH et al. 2012)		1, 2 and 3	x	↓		↑		↓	↓		↓
Ritz (Ritz B et al. 2014)		1 (weeks 0-19) 2 (weeks 19-29) 3 (weeks 29-37)						↓	x	x	
Iniguez(Iniguez et al. 2016)		1 (weeks 1-12) 2 (weeks 12-20) 3 (weeks 20-34 and 34-term)	x	x	↓ ψ	↓ ψ		↓ ψ	x	↓ ψ	↓ ψ
Carvalho(Carvalho M.A. et al. 2016)		1 (week 12) 2 (week 22) 3 (week 32)									x
Malmqvist(Malmqvist et al. 2017)		1, 2 and 3						x	↓	↓	↓
Clemens(Clemens et al. 2017)		Average across all trimesters	x	x	x			↓	x	x	
Handmaker (Handmaker NS et al. 2006)	Maternal alcohol	2 (<28 weeks)						↓	x	x	
Kfir (Kfir M et al. 2009)		2 (18 weeks)	x	↓	x	x		↓	x	x	x
Bakker (Bakker R et al.		1 (<18 weeks)	x	x	x	x		x	x	x	↑

2010)		2(18-25 weeks) 3 (>25 weeks)								
Snijder(Snijder CA et al. 2012b)(Occupational chemicals)	Other exposures	2 ("mid pregnancy")					↓	↓		↓
Snijder(Snijder CA et al. 2013)(Bisphenol A)		1, 2 and 3	x			x	x			x
Philippat(Philippat et al. 2014)			x	x	x	x	x	x	x	↓ψ
Harari(Harari et al. 2015)(Maternal plasma lithium)		1 or 2 or 3	x	x	x	x	x	x	x	x
Lopez-Espinosa(Lopez-Espinosa et al. 2015)(Flame retardants)		1 (weeks 10-13)	x	x	x	x	↓ψ	x	↓ψ	↓ψ
Botton(Botton et al. 2016)(Phthalates)		2 (week 26)		↓		x		↓		x
Casas(Casas et al. 2016)(Bisphenol A and Phthalates)		1 (12 weeks) 3 (32 weeks)	↓ψ**	x	x	x	x	↓ψ**	x	x
Lopez-Espinosa (Lopez_Espinosa et al. 2016) (organochlorine compounds)		1 (12 weeks)	x	x	x	x	x	↓ψ	x	↓ψ

* P=periconception, 1=first trimester, 2=second trimester, 3=third trimester. †limited to those whose mothers started folate supplementation before conception. §femur diameter and not femur length. ‡growth and not size reported. δ analysis limited to head circumference growth between the second and third trimester. ¥ see table 2 for more details of associations between individual exposures and fetal measurements. ψ second trimester details are growth between weeks 12-20 and third trimester details are for growth between weeks 20-34. **reduction in association with phthalates and not bisphenol A

Table two

Table 2. Details of studies where maternal ambient air exposures were linked to fetal ultrasound measurements. HC=head circumference, BPD=biparietal diameter, AC=abdominal circumference, EFW=estimated fetal weight. NO₂= nitrogen dioxide, CO=carbon monoxide, PM₁₀=particulates with diameter less than 10 microns, SO₂=sulphur dioxide.

	Number of mothers	Methodology for measuring ambient air exposure	Exposures measured and at what gestation	Units for comparing exposure	Association with fetal size/growth
Hansen (Hansen CA et al. 2008)	14,734	Monthly average of daily exposure from closest monitoring station	PM ₁₀ , Ozone, NO ₂ and SO ₂ over first four months	5 microg/m ³ for PM ₁₀ 5ppb for NO ₂ 8ppb for ozone 0.8ppb for SO ₂	1mm reduction in T2 BPD and SO ₂ during month 1 0.2mm reduction in T2 FL and PM ₁₀ during months 1 and 4 1mm reduction in T2 AC and ozone and PM ₁₀ during month 2 2mm reduction in T2 AC and SO ₂ during month 3 1mm reduction in T2 HC and PM ₁₀ during month 4
Slama (Slama R et al. 2009)	271	Non-smoking mothers carried diffusive air sampler	Benzene exposure at 27 weeks gestation	log transformed benzene exposure (microg/m ³)	1.5 mm reduction in T2 HC 2mm reduction in T3 HC
Aguilera(Aguilera I et al. 2010)	562	Land-use regression modelling using ambient measurements	NO ₂ and aromatic hydrocarbons (BTEX, including benzene, toluene) between weeks 1 and 32	microg/m ³	For all pregnancies 5% reduction in BPD growth between weeks 20-32 and BTEX weeks 1-12 For mothers who spent<2 hours a day out of the home, 6% reduction in HC growth between weeks 12-20 and 5% reduction in AC and EFW growth weeks 20-32 and NO ₂ weeks 1-12
Iniguez (Iniguez C et al. 2012)	818	Land-use regression modelling	NO ₂ between week 0 and delivery	microg/m ³	2-3% reduction in BPD growth weeks 20-32 and NO ₂ weeks 0-20 2% reduction in AC and EFW growth weeks 20-32 and NO ₂ weeks 12-20
van den Hooven(van den Hooven EH et al. 2012)	7772	Dispersion modelling	PM ₁₀ and NO ₂ throughout pregnancy	Quartiles and microg/m ³	0.1mm increase in CRL and current PM ₁₀ 0.2 mm reduction in third trimester HC and current PM ₁₀ 0.3g increase in second trimester EFW and current PM ₁₀ 0.1-0.2 mm reduction in second trimester FL and in third trimester FL and HC and current NO ₂

Ritz (Ritz B et al. 2014)	500	Closest monitor methods for CO; NO ₂ ; O ₃ ; PM ₁₀ . Dispersion modelling and land-use regression for NO _x	Mean daily NO ₂ , nitrogen oxides (NO _x), CO, ozone, PM ₁₀ between weeks 0-19, 19-29 and 29-37.	Quartiles of exposure	0.2-1 reduction in HC at 37 weeks per quartile increase in PM ₁₀ between weeks 27-29. ~1mm reduction in HC at 37 weeks per quartile increase in NO ₂ between weeks 27-39
Iniguez(Iniguez et al. 2016)	2478	Land-use regression modelling	NO ₂ for weeks 0-12, 12-20, 20-34 and 34-delivery	10 microg/m ³	2-3% reduced BPD and AC growth weeks 20-34 and exposure <20 weeks (not FL growth)
Carvalho(Carvalho M.A. et al. 2016)	366	Passive personal samplers	NO ₂ and Ozone for 7-18 days before first, second and third trimester ultrasound scan	Log transformed microg/m ³	No association with EFW in third trimester. Ozone exposure was associated with umbilical artery characteristics
Malmqvist(Malmqvist et al. 2017)	48,000	Dispersion modelling 500m or 100m grid cells	Nitrogen oxides for each trimester	10 microg/m ³	0.1mm reduction T2 AD and FL (not T2 BPD) 0.16 z score reduction T2 EFW 0.5 mm reduction head circumference 0.7 z score change in growth during T3
Clemens(Clemens et al. 2017)	13,755	Dispersion modelling 1km cells	Average NO ₂ , PM ₁₀ , PM _{2.5} throughout pregnancy	Quartiles of exposure	0.3 z score reduction in T3 BPD in highest versus lowest PM ₁₀ exposure quartile

Figure one
[Click here to download high resolution image](#)




Figure two
[Click here to download high resolution image](#)

Author	Exposure	Recruitment strategy	Blinding	Exposure assessment	Confounding	Incomplete outcome	Selective reporting	Conflict of interest	Other bias
Bergen	Diet (Homocysteine)								
Bouwland-Both	Diet								
Carlsen	Diet (Fatty acids)								
Drouillet	Fish pre preg								
Drouillet-Pinard	Mercury pre preg								
Heppe	Fish								
Heppe	Milk								
Ioannou	Vitamin D								
Karateke	Fasting								
Mahon	Vitamin D								
Steenweg	Folic acid								
Timmermans	Folic acid								
Timmermans	Mediter diet								
Turner	Vitamin E								
Young	Vitamin D								
Aguilera	Air pollution								
Carvalho	Air pollution								
Clemens	Air pollution								
Hansen	Air pollution								
Iniguez	Air pollution								
Iniguez	NO2								
Malmqvist	Air pollution								
Ritz	Air pollution								
Slama	Benzene								
van den Hooven	Air pollution								
Bakker	Alcohol								
Handmaker	Alcohol								
Kfir	Alcohol								
Botton	Phthalate								
Casas	Bisphenol A								
Harari	Lithium								
Lopez-Espinosa	Flame retardants								
Phillipat	Phenols in Boys								
Snijder	Bisphenol A								
Snijder	Occupational chemicals								
Lopez-Espinosa	Organochlorine compounds								

Risk of bias


Low

Probably low

Probably high

High

Figure three
[Click here to download high resolution image](#)



Supplementary Material

[Click here to download Supplementary Material: Supplement for resubmission .docx](#)

A systematic review of associations between maternal exposures during pregnancy other than smoking and antenatal fetal measurements

Ivory Huang^a, Diane Mak^a, Prudence Cheung^a, Miriam Abraham^a, Tom Clemens^b,
Steve Turner^a

^aChild Health, University of Aberdeen, Aberdeen, UK

^bSchool of Geosciences, University of Edinburgh, UK

Correspondence to: Prof Turner, Child Health, Royal Aberdeen Children's Hospital, Aberdeen, AB25 2ZG, UK. Tel +44 1224 438470. s.w.turner@abdn.ac.uk

Declarations of interest: None

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors

Author contributions: IH and MA designed the search methodology and contributed to the first draft of the manuscript. DM and PC completed the assessment of the identified literature. TC contributed to the interpretation of the air pollution literature. ST conceived the study, was involved in the search methodology, assessment of the literature and contributed to the first draft of the manuscript. All authors made meaningful contributions to the submitted manuscript.